

## Original Article

## Seroprevalence of Hepatitis E Virus Infection among Patients with Acute Hepatitis Symptoms in Ahvaz, Iran

Mehdi Parsa Nahad<sup>1,2</sup>Ph.D., Akbar Bavi<sup>1,2</sup>M.Sc., Milad Zandi<sup>3</sup>Ph.D.  
 Mehrab Nejati<sup>4</sup>Ph.D., Mehdi Rabie Rudsari<sup>5</sup>M.Sc., Mohammad Amin  
 Dehghani<sup>6</sup>M.Sc., Hadis Kiani<sup>1,2</sup>M.Sc., Ali Ramezani<sup>1,2\*</sup> Ph.D.  
 Sara Khosravi Seftajani<sup>7</sup>M.Sc., Nahid Omidi<sup>1,2</sup>M.Sc., Hojatolla  
 Nikravesh<sup>6</sup>Ph.D., Seyed Arash Moradpour<sup>8</sup>Pharm.D.

<sup>1</sup>Health Research Institute, Infectious and Tropical Diseases Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

<sup>2</sup>Department of Virology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

<sup>3</sup>Department of Virology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

<sup>4</sup>Department of Microbiology, Shahre Babak Azad University, Shahre Babak, Iran.

<sup>5</sup>Department of Microbiology, Golestan University of Medical Sciences, Gorgan, Iran.

<sup>6</sup>Department of Pharmacology and Toxicology, School of Pharmacy and Toxicology Research Center, Jundishapur University of Medical Sciences, Ahvaz, Iran.

<sup>7</sup>Department of Medical Microbiology, Faculty of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran.

<sup>8</sup>Department of Pharmaceutics, School of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

### ABSTRACT

#### Article history

Received 29 Jun 2017

Accepted 13 Dec 2017

Available online 18 Mar 2018

#### Key words

HEV

Endemic

Fulminant

Seroprevalence

**Background and Aims:** Hepatitis E virus (HEV) is considered as one of the common causes of acute viral hepatitis in the world, particularly in the developing countries. It also causes fulminant hepatitis in pregnant women. Although HEV infection in Iran is endemic, there are numerous reports of epidemics caused by this infection in this part of the world. This study aimed to determine seroprevalence of HEV among patients with acute hepatitis symptoms in Ahvaz, Iran.

**Materials and Methods:** In this cross-sectional study patients were selected based on laboratory findings related to acute hepatitis including elevated serum bilirubin levels, markedly elevated levels of liver enzymes such as alanine aminotransferase, aspartate aminotransferase and mild increases in alkaline phosphatase activity. The plasma samples were tested for IgM/IgG Hepatitis E antibody (anti-HEV) using enzyme-linked immunosorbent assay.

**Results:** Out of 241 study patients, 125 (51.9%) were male and 116 (48.1%) female. The mean age $\pm$ SD was 31.53 $\pm$ 17.17 years. The results indicated 27.4% prevalence of anti-HEV antibodies among patients with acute hepatitis in Ahvaz.

**Conclusions:** In regard with the findings, HEV can play an important role in incidence of hepatitis. The study needs to be conducted in high-risk groups and in general populatios.

**\*Corresponding Author:** Health Research Institute, Infectious and Tropical Diseases Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. **Tel:** +986133738392, **Email:** Ramezani.al@ajums.ac.ir

## Introduction

Hepatitis E virus (HEV) is an icosahedral non-enveloped, single-stranded positive-sense RNA virus in the genus of Hepevirus from the family Hepeviridae [1]. HEV is considered as one of the most common causes of acute viral hepatitis over the world particularly in the developing countries. This is a major cause of fulminant hepatitis in pregnant women [2, 3]. The viral particle is relatively stable under environmental conditions like sewage water [4]. Therefore, the oral-fecal route, like contaminated food or water, is a predominant route of HEV transmission to others. In rare cases, HEV can be transmitted zoonotically and by blood transfusion [5]. However, there is evidence of vertical transmission of hepatitis E virus infection from mother to fetus seemingly representing the ability of the virus to cause congenital infections [6]. HEV infection is regarded as a significant public health problem especially in the developing countries where large outbreaks and epidemics of HEV infections happen as a result of poor personnel hygiene, inadequate sanitation, and an unsafe drinking water supply [7]. Several studies have also reported the high seroprevalence of HEV infection in the developed countries [8]. The clinical symptoms and laboratory findings of HEV infection are similar to other hepatotropic viruses which cause acute hepatitis [9]. Pregnant women with HEV infection in the third trimester are at an increased risk of fulminant hepatic failure and maternal death [10]. One-third of the world population is infected with HEV virus [11]. Each year, 20 million new

infections and 3.3 million acute infections as well as 56,600 deaths occur in the world [12, 13]. In general populations, the mortality rate is 1-2% [11]. However, in pregnant women and in individuals with pre-existing liver disorders it may amount to 10-25% and over 75%, respectively [14]. Iran is considered as an endemic area of HEV infection [15]. The prevalence of HEV infection in Iran is very different due to the variations in public health and hygiene levels, risk factors and transmission routes in different areas and groups [16]. According to the studies conducted in different regions of Iran, the HEV prevalence is reported to be 1.1-14.2% in public population [17], 4.5-14.3% in blood donors [18], 6.3-8.3% in patients of hemodialysis [19], 27.5% in chronic liver disease [20], and 10-16.4% in HIV-infected patients [3]. However, Iran is the endemic area of HEV infection and studies carried out in this regard are not adequate [15]. Therefore, this study was performed to evaluate the seroprevalence of HEV in patients with acute hepatitis symptoms.

## Materials and Methods

### Study population

In this cross-sectional study, 241 patients including 125 (51.9%) male and 116 (48.1%) female (mean age $\pm$ SD: 31.53 $\pm$ 17.17 years) with acute hepatitis symptoms such as elevated liver enzymes levels such as increased serum alanine aminotransferase, aspartate aminotransferase and bilirubin as well as clinical symptoms such as fever, fatigue,

malaise, anorexia, jaundice and pruritus were enrolled during December 2013 to August 2016 in the city of Ahvaz. This project was approved by the Ethic Committee of Ahvaz Jundishapur university of medical sciences. In addition, the consent form was obtained from all the patients.

#### **Detection of antibodies against HEV**

The peripheral venous blood was collected from each patient. Plasma samples were separated from whole blood. Samples were tested for IgM/IgG Hepatitis E antibody (anti-HEV) using enzyme-linked immunosorbent assay (ELISA) according to the manufacturer's instruction (DIA.PRO HEV Ab ELISA kit).

#### **Statistical analyses**

Statistical analyses were performed using Chi-square and SPSS 21 Package program (SPSS Inc., Chicago, IL, USA). The p-values of lower than 0.05 were regarded as statistically significant. The proportions of patients which are categorized by sex, age and seasons was compared by ANOVA test and Fisher's exact statistics. Descriptive statistics were used for data analysis. The results were presented as frequencies or percentage.

## **Results**

In terms of age, the patients were divided into six groups comprising less than 15 yrs, 16-25 yrs, 26-35 yrs, 36-45 yrs, 46-55 yrs and more than 56 yr. Those with 42/241 (17.4%), 59/241 (24.5%), 56/241 (23.2%), 36/241 (14.9%),

18/241 (7.5%) and 30/241 (12.4%) constructed groups 1 to 6 respectively (Fig. 1). In terms of season, 72/241 (29.9%) patients were in the spring 54/241 (22.4%) in the summer, 45/241 (18.7%) were in the fall, and 70/241 (29%) patients in the winter (Fig. 2). Table 1 presents a summary of characteristics of the study of all the patients. Of the 241 patients, 66/241 (27.4%) were shown to be positive for anti-HEV total antibodies while 175/241 (72.6%) were negative. The overall anti-HEV total antibodies prevalence rate was 27.4% among patients with acute hepatitis. Based on the gender and season, 30/125 in the male group and 36/116 in the female group were positive for anti-HEV total antibodies.

Totally, 21/72 in the spring, 15/54 in the summer, 15/45 in the fall, and 15/70 in the winter were shown to be positive for anti-HEV total antibodies. Although, the seroprevalence was higher in the spring season and female groups; but HEV seropositivity was not statistically significant in terms of gender ( $p=0.223$ ) and season ( $p=0.542$ ). Nevertheless, there was a statistically significant difference in anti-HEV total antibodies seroprevalence rate between the patients grouped based on age ( $p<0.01$ ). Seroprevalence of (HEV) antibody increased with age from 7.1% (3/42) in group 1 to 76.7% (23/30) in group 6, so that the highest rate of anti-HEV seroprevalence was identified in group 6 (Table 1).




Fig .1. Prevalence of anti-HEV IgM/IgG antibody in categorized by age




Fig. 2. Prevalence of anti-HEV IgM/IgG antibody according to four seasons

**Table 1.** Anti-HEV total antibodies seroprevalence rate in studied samples

	Participants (n=241)	Anti-HEV positive (n=66)	Anti-HEV Negative (n=175)	P-value
<b>Gender</b>				0.223
<b>Male</b>	125 (51.9%)	30 (45.5%)	95 (54.3%)	
<b>Female</b>	116 (48.1%)	36 (54.5%)	80 (45.7%)	
<b>Season</b>				0.542
<b>Spring</b>	72 (29.9%)	21 (29.2%)	51 (70.8%)	
<b>Summer</b>	54 (22.4%)	15 (27.8%)	39 (72.2%)	
<b>Fall</b>	45 (18.7%)	15 (33.3%)	30(66.7%)	
<b>Winter</b>	70 (29.0%)	15 (21.4%)	55 (78.6%)	
<b>Age groups (yr)</b>				<0.01
<b>≤15</b>	42 (17.4%)	3 (7.1%)	39 (92.9%)	
<b>16-25</b>	59 (24.5%)	9 (15.3%)	50 (84.7%)	
<b>26-35</b>	56 (23.2%)	8 (14.3%)	48 (85.7%)	
<b>36-45</b>	36 (14.9%)	13 (36.1%)	23 (63.9%)	
<b>46-55</b>	18 (7.5%)	10 (55.6%)	8 (44.4%)	
<b>≥56</b>	30 (12.4%)	23 (76.7%)	7 (23.3%)	

## Discussion

Hepatitis E infection is a global public health problem causing large outbreaks of acute hepatitis in the developing countries especially in the Southeast and Central Asia, the Middle East, northern and western parts of Africa as well as sporadic cases of the infection also in the developed countries [21]. There is wide variation of HEV prevalence in different areas of the world [22], however the HEV prevalence is greater in tropical and subtropical regions [23]. Since the overall prevalence rate of HEV in Iranian general population is more than 5%, Iran is considered as an endemic zone for HEV disease [15, 24]. Previous studies have reported variations in the epidemiology and prevalence of hepatitis E in different regions of Iran. These variations are due to differences in the life styles, status

of public health, risk factors, and routes of transmission in different groups and geographical areas [25]. Taremi et al. reported HEV seroprevalence rate of 9.3% among general population in Nahavand Province in Iran [26]. Raofi et al. indicated that HEV seroprevalence rate in general population of Khorramabad city is 7.8% [27]. In another study by Mohebbi et al. the prevalence of HEV was reported to be 9.3% in general population of Tehran [28]. Also Farshadpour et al. reported HEV seroprevalence rate leveling 46.1% among adults in South-West of Iran [29]. In Arabzadeh et al.'s study, HEV prevalence was 7.7% among blood donors in kerman [30]. Therefore, the available epidemiological studies in different regions of Iran report the highest rate of HEV prevalence

related to Ahvaz, the center of Khozestan province. Perhaps one of the main reasons for the high prevalence of this virus in Ahvaz is the pollution of Karun river water as the only source of drinking water in the region. This results from the fact that in many cases, household sewage, industrial wastewater, and wastewater from agriculture and hospitals enter the Karoon river. There are also reports of some epidemiologic studies performed on HEV prevalence among the general populations of other countries: 28% in Hong Kong [31], about 3.4% in Japan blood donors [32], 9% in Vietnam [33], 1.1% in Spain [34], and 3.6% in the general population of China [35]. The present probe investigated the HEV seroprevalence among the patients with acute hepatitis in Ahvaz city and found anti-HEV total antibodies to be 27.4%. In this study, the HEV seroprevalence rate significantly increased with age; from 7.1% in people aged below 26 years to 76.7% in individuals aged more than 56 years. The current study, in line with some other studies, indicated a statistically significant difference between age and the prevalence of the disease increasing with age [36, 37]. In accordance with the results of the previous studies [38-40], our findings showed that the presence of anti-HEV total antibodies is not associated with gender. Outbreaks of hepatitis E in the tropical developing countries mainly occurs during the rainy season due to floods when sewage water gains access to open water reservoirs, as well as the overcrowding factor [41]. However our results showed a statistically insignificant difference in anti-HEV total antibodies

seroprevalence rate between seasons, In other words, the presence of anti-HEV total antibodies is not associated with a particular season. Given that there is no effective vaccine to prevent infection and also due to the risk of progression to advanced liver disease including fulminant hepatitis or even chronic infection and cirrhosis with a high rate of mortality in high-risk groups like pregnant women, organ transplant recipient patients, and immunosuppressed patients, research on HEV infection epidemiology is very critical so as to employ preventive measures in the public health services of the country [13, 42-44].

## Conclusions

Overall, the results of this study indicate that HEV can play an important role in the incidence of acute hepatitis. Therefore, it seems that further epidemiological investigations are needed to evaluate the association and role of the virus in the development of chronic hepatitis. Regarding the high prevalence of this disease, it seems that improving the quality of drinking water in Ahvaz should be considered as one of the most vital priorities. On the other hand, in the context of research, screening for this disease appears to be necessary in high-risk groups and at-risk populations.

## Conflict of Interest

There is no conflict of interest in this article.

## Acknowledgements

This study was done in virology department of Ahvaz Jundishapur University of Medical Sciences. The authors are very thankful to all patients for providing samples for scientific research.

## References

- [1]. Meng XJ. Hepatitis E virus: animal reservoirs and zoonotic risk. *Vet Microbiol.* 2010; 140(3-4): 256-65.
- [2]. Navaneethan U, Al Mohajer M, Shata MT. Hepatitis E and pregnancy: understanding the pathogenesis. *Liver Int.* 2008; 28(9): 1190-199.
- [3]. Joulaei H, Rudgari O, Motazedian N, Gorji-Makhsous S. Hepatitis E virus seroprevalence in HIV positive individuals in Shiraz, Southern Iran. *Iran J Microbiol.* 2015; 7(2): 103-8.
- [4]. Hughes JM, Wilson ME, Teshale EH, Hu DJ, Holmberg SD. The two faces of hepatitis E virus. *Clinic Infect Dis.* 2010; 51(3): 328-34.
- [5]. Yugo DM, Meng XJ. Hepatitis E virus: foodborne, waterborne and zoonotic transmission. *Int J Environ Res Public Health.* 2013; 10(10): 4507-533.
- [6]. Zaki ME, Aal AA, Badawy A, El-Deeb DR, El-Kheir NY. Clinicolaboratory study of mother-to-neonate transmission of hepatitis e virus in egypt. *Am J Clinic Pathol.* 2013; 140(5): 721-26.
- [7]. Kumar S, Subhadra S, Singh B, Panda BK. Hepatitis E virus: The current scenario. *Int J Infect Dis.* 2013 Apr 30; 17(4): e228-33.
- [8]. Emerson SU, Purcell RH. Hepatitis E virus. *Rev Med Virol.* 2003; 13(3): 145-54.
- [9]. Shrestha AC, Faddy HM, Flower RL, Seed CR, Keller AJ. Hepatitis E virus: do locally acquired infections in Australia necessitate laboratory testing in acute hepatitis patients with no overseas travel history? *Pathology* 2015; 47(2): 97-100.
- [10]. Chaudhry SA, Verma N, Koren G. Hepatitis E infection during pregnancy. *Can Fam Physician.* 2015; 61(7): 607-608.
- [11]. Kim JH, Nelson KE, Panzner U, Kasture Y, Labrique AB, Wierzb TF. A systematic review of the epidemiology of hepatitis E virus in Africa. *BMC Infect Dis.* 2014; 14: 308.
- [12]. Bazerbachi F, Haffar S, Garg SK, Lake JR. Extra-hepatic manifestations associated with hepatitis E virus infection: a comprehensive review of the literature. *Gastroenterol Rep (Oxf).* 2016; 4(1): 1-15.
- [13]. Sridhar S, Lau SK, Woo PC. Hepatitis E: A disease of reemerging importance. *J Formos Med Assoc.* 2015; 114(8): 681-90.
- [14]. Marano G, Vaglio S, Pupella S, Facco G, Bianchi M, Calizzani G, et al. Hepatitis E: an old infection with new implications. *Blood Transfus.* 2015; 13(1): 6-17.
- [15]. Sepanlou SG, Rezvan H, Amini-Kafiabad S, Dayhim MR, Merat S. A population-based seroepidemiological study on hepatitis E virus in Iran. *Middle East J Dig Dis.* 2010; 2(2): 97-103.
- [16]. Seyed Mohammad A, Fatameh A, Mohammad Reza G. Seroepidemiological Study of Hepatitis E Virus in Drug Addicts in Ahvaz, Southern Iran: 2005-2006. *Hepat Mon.* 2008(4): 263-66.
- [17]. Asaei S, Ziyaeyan M, Moeini M, Jamalidoust M, Behzadi MA. Seroprevalence of hepatitis A and E virus infections among healthy population in Shiraz, Southern Iran. *Jundishapur J Microbiol.* 2015; 8(7): e19311.
- [18]. Taremi M, Gachkar L, Mahmoud Arabi S, Kheradpezhoh M, Khoshbaten M. Prevalence of antibodies to hepatitis E virus among male blood donors in Tabriz. Islamic Republic of Iran. *East Mediterr Health J.* 2007; 13(1): 98-102.
- [19]. Hosseini-Moghaddam SM, Zarei A, Alavian SM, Mansouri M. Hepatitis E virus infection: a general review with a focus on hemodialysis and kidney transplant patients. *Am J Nephrol.* 2010; 31(5): 398-407.
- [20]. Keyvani H, Shamabadi MS, Najafifard S, Hajibeigi B, Fallahian F, Alavian SM. Seroprevalence of anti-HEV and HEV RNA among volunteer blood donors and patients with Hepatitis B and C in Iran. *Bangladesh Liver Journal* 2009; 1(1): 34-7.
- [21]. Ghezdasht SA, Miri R, Hedayatimoghadam M, Shamsian A, Bidkhorji H, Fathimoghadam F, et al. Population movement and virus spreading: HEV spreading in a Pilgrimage city, Mashhad in northeast Iran; an example. *Hepat Mon.* 2013; 13(8): e10255.
- [22]. Assarehzadegan MA, Shakerinejad G, Amini A, Rezaee SR. Seroprevalence of hepatitis E virus in blood donors in Khuzestan province, southwest Iran. *Int J Infect Dis.* 2008; 12(4): 387-90.
- [23]. Behzadifar M, Lankarani KB, Abdi S, Mirghaed MT, Beyranvand G, Keshavarzi A, et al. Seroprevalence of Hepatitis E Virus in Iran: A Systematic Review and Meta-analysis. *Middle East J Dig Dis.* 2016; 8(3): 189-200.
- [24]. Alavian SM, Fallahian F, Lankarani KB. Epidemiology of hepatitis E in Iran and Pakistan. *Hepat Mon.* 2009; 9(1): 60-65.
- [25]. Taherkhani R, Farshadpour F. Epidemiology of hepatitis E virus in Iran. *World J gastroenterol.* 2016; 22(22): 5143-153.
- [26]. Taremi M, Alizadeh AH, Ardalan A, Ansari S, Zali MR. Seroprevalence of hepatitis E in Nahavand, Islamic Republic of Iran: a population-based study. *East Mediterr Health J.* 2008; 14(1): 157-62.

- [27].Raofi R, Nazer MR, Pournia Y. Seroepidemiology of hepatitis E virus in Western Iran. *Braz J Infect Dis.* 2012; 16(3): 302-303.
- [28].Mohebbi SR, Rostami Nejad M, Tahaei SM, Pourhoseingholi MA, Habibi M, Azimzadeh P, et al. Seroepidemiology of hepatitis A and E virus infections in Tehran, Iran: a population based study. *Transactions of the Royal Society of Tropical Medicine and Hygiene.* 2012;106(9):528-31.
- [29].Farshadpour F, Taherkhani R, Makvandi M. Prevalence of hepatitis E virus among adults in south-west of Iran. *Hepat Res Treat.* 2015; 2015.
- [30].Arabzadeh SA, Zahedi M, Molaei H, Aghaei AA. Seroepidemiology of Anti-HEV IgG in Healthy Men Blood Donors in Kerman, 2007-2008. *Iranian J virol.* 2010; 3(1): 30-2.
- [31].Chiu DM, Chan MC, Yeung A, Ngai KL, Chan PK. Seroprevalence of hepatitis E virus in Hong Kong, 2008-2009. *J Med Vir.* 2013; 85(3): 459-61.
- [32].Takeda H, Matsubayashi K, Sakata H, Sato S, Kato T, Hino S, et al. A nationwide survey for prevalence of hepatitis E virus antibody in qualified blood donors in Japan. *Vox Sang.* 2010; 99(4): 307-13.
- [33].Hau CH, Hien TT, Tien NT, Khiem HB, Sac PK, Nhung VT, et al. Prevalence of enteric hepatitis A and E viruses in the Mekong River delta region of Vietnam. *Am J Trop Med Hyg.* 1999; 60(2): 277-80.
- [34].Fogeda M, Avellón A, Echevarria JM. Prevalence of specific antibody to hepatitis E virus in the general population of the community of Madrid, Spain. *J Med Vir.* 2012; 84(1): 71-4.
- [35].Lu B, Zhao HL, Tian RG, Li HY, Zhan J, Wang XY, et al. Antibody detection of hepatitis E virus in human population of different national in China. *Zhonghua shi yan he lin chuang bing du xue za zhi= Zhonghua shiyan he linchuang bingduxue zazhi= Chinese J Experiment Clin Vir.* 2012; 26(1): 46-7.
- [36].Faber MS, Wenzel JJ, Jilg W, Thamm M, Höhle M, Stark K. Hepatitis E virus seroprevalence among adults, Germany. *Emerging Infect Dis.* 2012; 18(10): 1654-657.
- [37].Tadesse E, Metwally L, Alaa EL. High prevalence of anti-hepatitis E virus among Egyptian blood donors. *J General Mol Vir.* 2013; 5(1): 9-13.
- [38].Kaufmann A, Kenfak-Foguena A, André C, Canellini G, Bürgisser P, Moradpour D, et al. Hepatitis E virus seroprevalence among blood donors in southwest Switzerland. *PloS one* 2011; 6(6): e21150.
- [39].Ehteram H, Ramezani A, Eslamifar A, Sofian M, Banifazl M, Ghassemi S, et al. Seroprevalence of Hepatitis E Virus infection among volunteer blood donors in central province of Iran in 2012. *Iran J Microbiol.* 2013; 5(2): 172-76.
- [40].Ataei B, Nokhodian Z, Javadi AA, Kassaian N, Shoaie P, Farajzadegan Z, et al. Hepatitis E virus in Isfahan Province: a population-based study. *Int J Infect Dis.* 2009; 13(1): 67-71.
- [41].Alavian SM. A Look at the Past History of Hepatitis E in Haiti: Should it be a Warning sign during the Current Crisis? *Hepatitis monthly.* 2010; 10(1): 9.
- [42].Pérez-Gracia MT, García M, Suay B, Mateos-Lindemann ML. Current knowledge on hepatitis E. *J Clin Transl Hepatol.* 2015; 3(2): 117-26.
- [43].Bajpai M, Gupta E. Transfusion-transmitted hepatitis E: is screening warranted? *Indian J Med Microbiol.* 2011; 29(4): 353-58.
- [44].Kamar N, Dalton HR, Abravanel F, Izopet J. Hepatitis E virus infection. *Clinic Microbiol Rev.* 2014; 27(1): 116-38.