

Original Article

The Protective Effect of Silymarin on Lipopolysaccharide-Induced Liver Toxicity in Male Wistar Rat

Fahimeh Nourbakhsh ¹Ph.D., Samaneh Borooni ²M.Sc.
Elaheh Tajbakhsh ^{3*}Ph.D.

¹Medical Toxicology Research Centre, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

²Department of Biology, Nourdanesh Institute of Higher Education, Meymeh, Esfahan, Iran.

³Department of Microbiology, Faculty of Basic Sciences, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran.

ABSTRACT

Article history

Received 28 Jan 2019

Accepted 2 Jul 2019

Available online 10 Dec 2019

Key words

Apoptosis

Liver inflammation

LPS

Oxidative stress

Silymarin

Background and Aims: Lipopolysaccharide (LPS) triggers production of reactive oxygen species and inflammatory cytokines. Nowadays *Silybum marianum* has been shown to treat liver and gall bladder disorders, especially to protect the liver against poisoning from various toxins. Therefore, we decided to evaluate the protective effect of silymarin on LPS-induced liver toxicity in male Wistar rat.

Materials and Methods: Totally, 40 male Wistar rats were divided into 4 groups (n=10 in each): The animals were treated with silymarin for two weeks before the biochemical tests. Apoptosis was assessed by evaluating the amount of proteins in liver tissues by western blotting.

Results: LPS induced hepatotoxicity as evidenced by histopathological damages and biochemical abnormalities. Data showed that malondialdehyde level significantly increases in the liver of LPS-treated rats. Destructive effects of LPS on histopathological and biochemical parameters were improved. LPS also increased expression of Bax/Bcl₂ ratio and activation of caspase 3, caspase 8 and caspase 9. Western blot analysis showed silymarin treatment inhibiting apoptosis stimulated by LPS in the liver (p<0.001).

Conclusions: The results of this research demonstrated that silymarin can exert protective effects against toxic effects of LPS in rat liver. Anti-inflammatory drug can play a protective role in attenuating the liver inflammation induced by LPS injection.

***Corresponding Author:** Departments of Microbiology, Faculty of Basic Sciences, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran. **Tel/Fax:** +9891322399694, **Email:** ee_tajbakhsh@yahoo.com

Introduction

The liver is an important organ which has a central role in metabolic homeostasis and is responsible for the metabolism, synthesis, storage and redistribution of nutrients, carbohydrates, fats and vitamins [1]. The liver plays a key role in immunological homeostasis and metabolism. This organ is also known to be an important place for generation of free radicals. In addition, free radical production can occur if inflammation is caused by or exposed to toxins [2]. Liver enzymes, including diamine oxidase, aldehyde dehydrogenase, tryptophan dual oxidase, liver dehydrogenase and the cytochrome P450 enzyme system, induce oxidation and production of uncoupling molecules [3, 4]. A high number of free radicals in the liver would be affected by endogenous or exogenous antioxidants and drugs such as acetaminophen. Molecular oxygen has a unique electronic structure that can accept a total of four electrons, and frequently evolve into free radicals formation. Oxygen free radicals, more generally known as reactive oxygen species (ROS), along with reactive nitrogen species represent the most important group of radical species generated in living system. Liver inflammation produces excessive oxygen free radicals to attack host cells, leading to cell damage [5]. Peroxidation of DNA plays an important role in pathogenesis of mutations in the liver cells. The activity of superoxide dismutase, and glutathione peroxidase in chronic liver cirrhosis and hepatitis is significantly low, which is negatively correlated with serum alanine aminotransferase (ALT)

level [6]. Lipopolysaccharide (LPS) is a major constituent of the outer membrane of gram-negative bacteria, and the immune system is constantly exposed to low levels of LPS through low-grade bacterial infections [7, 8]. LPS recognition and signaling responses are keys to eliminating invading pathogens. LPS-induced activation of macrophages induces production of bioactive lipids, ROS and in particular, inflammatory cytokines. Although, the LPS response is critical to fighting and clearing bacterial infections, LPS also mediates deleterious host reactions. In addition, LPS can initiate extra production of inflammatory mediators, leading to potentially lethal systemic diseases, for example, septic shock and tissue damage. Inflammation can induce oxidative damage by generating oxidative agents such as the superoxide anion and nitric oxide (NO) [9, 10]. Inducible NO synthase (iNOS) produce NO and NO reacts with superoxide and generate peroxynitrite, particularly in immune cells [11]. Additionally, NO synthesizes oxidize the terminal guanidino nitrogen of L-arginine, producing NO and citrulline. Various extracellular stimuli activate nuclear factor kappa-light-chain-enhancer of activated B-cells, an inducible transcription factor. Stimuli include cytokines, LPS and oxidative stress that can cause chronic liver impairment with activation of hepatic stellate cells, recruitment of inflammatory cells, activation of infiltrating immunocompetent cells and initiation of oxidative stress cascade [12]. Silymarin, a standardized extract obtained from seeds of

Silybum marianum (*S. marianum*), is widely used in treatment of liver diseases of varying origins. Seeds of *S. marianum* have been shown to treat liver and gall bladder disorders, including hepatitis, cirrhosis and jaundice and to protect the liver poison from chemicals, environmental toxins, snake bites, insect stings, mushroom poisoning and alcohols. Due to its proven hepatoprotective and antioxidant properties, silymarin is being used as a standard agent for hepatoprotective effects induced by LPS [13, 14].

The emergence of silymarin as a natural remedy for liver diseases, coupled with its entry into national institutes of health clinical trial, signifies its hepatoprotective potential. Silymarin is noted for its ability to interfere with apoptotic signaling while acting as an antioxidant. Various *in vivo* studies were designed to explore the hepatotoxic potential of Doxorubicin, the well-known cardiotoxin, and in particular the possibility of pre-exposures to silymarin to prevent hepatotoxicity by reducing Dox-induced free radical mediated oxidative stress, by modulating expression of apoptotic signaling proteins like Bcl-xL, and by minimizing liver cell death occurring by apoptosis or necrosis [15]. In this way, we decided to evaluate the protective effect of silymarin on LPS-induced liver toxicity in male Wistar rat.

Materials and Methods

Animals and drugs

Totally, 40 male Wistar rats being 12 weeks old and 240±10 g weight were purchased from an animal house. The animals were kept in

standard conditions (temperature was 22±2°C and 12 h light/dark cycle). The rats had free access to food and water. The animals were divided into 4 groups (n=10 in each):

- 1) Control: received 1 ml/kg of saline instead of LPS or silymarin
- 2) LPS (daily injection): received 2 mg/kg LPS intraperitoneally with saline instead of silymarin (orally pretreatment in corn oil)
- 3) LPS- silymarin: received 10 mg/kg silymarin before 2 mg/kg LPS
- 4) LPS- silymarin: received 20 mg/kg silymarin before 2 mg/kg LPS

They were injected intraperitoneally with saline instead of LPS and silymarin for two weeks before the biochemical tests. LPS was purchased from Sigma (Sigma Chemical Co). silymarin was kindly provided by Daroupakhsh Company in Iran. In the final stage, urethane was injected to induce a deep anesthesia [16]. Then the livers were removed as liver toxicity index to evaluate the level of oxidative stress including malondialdehyde, catalase and liver enzymes including ALT and aspartate aminotransferase (AST). The Animal Care and Use Committee, and also the National Institutes of Health Guidelines for the Care and Use of Laboratory Animals approved this experimental protocol [17].

Determination of malondialdehyde and catalase activity

Malondialdehyde levels, as an index of lipid peroxidation, were measured in the liver tissues. malondialdehyde reacts with thiobarbituric acid as a thiobarbituric acid reactive substance and produces a red colored complex which has a peak absorbance at 535 nm. two ml

thiobarbituric acid/ trichloroacetic acid/ hydrochloric acid reagent was added to 1 ml homogenate and the solution was incubated in a boiling water bath for 40 min. After cooling, the whole solutions were centrifuged (1000 g for 10 min). The absorbance of supernatant was measured at 535 nm. The malondialdehyde concentration was calculated as follows: $C(m) = \text{Absorbance} / (1.65 \times 10^5)$ [18, 19].

Catalase activity was estimated using the method of Azizi et al. [12]. The principle of the assay is based on determination of the rate constant of hydrogen peroxide decomposition. By measuring the decrease in absorbance at 240 nm per minute, the rate constant of the enzyme was determined. Activities were expressed as k (rate constant) per liter. Catalase is a very important enzyme in protecting the cell from oxidative damage by ROS [20]. Liver tissues Interleukin (IL)-6 contents determination was performed with a specific rat enzyme-linked immunosorbent assay kit (bioscience Co, San Diego, CA, USA) according to Videla et al [19].

Biochemical blood evaluation

Serum levels of some liver enzymes like ALT and AST were assessed by using commercial colorimetric kits (Hitachi, Japan) [21, 22].

Western blot analysis

For western blot analysis, protein was extracted from the liver tissues for Bax, Bcl₂, caspase-3, caspase-8 and caspase-9. The total protein content was determined using Bradford protein assay kit (Bio-Rad). Levels of Bax, Bcl₂, caspase-3, caspase-8, caspase-9 and β-actin were measured by immune blotting analysis [23]. Briefly, equal amounts of protein

extracts (50 μg) were mixed with loading buffer and heated for 8 min at 95°C, loaded to sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) gel and separated by SDS-PAGE on a 12% gel. After electrophoresis, proteins were transferred to polyvinylidene fluoride membrane [24, 25]. The blots were incubated in blocking buffer tris buffered saline with tween 20 for 2 hr at room temperature. The primary antibodies were rabbit monoclonal anti-serum against Bcl₂, caspase-3, caspase-8, caspase-9, Bax and mouse monoclonal anti serum against β-actin (Cell Signaling). Finally, the membranes were incubated with the corresponding secondary antibody. Protein bands were visualized using an enhanced chemiluminescence (Pierce ECL Western blotting substrate) and Alliance gel doc (Alliance 4.7 Gel doc, UK). UV Tec software (UK) was used to semi-quantify protein bands. All protein bands were normalized against β-actin protein in order to achieve the final results.

Statistical analysis

All data were expressed as means ± SEM. The data were compared by one way ANOVA followed by tukey's post hoc comparisons test by prism 5. Differences were considered statistically significant when $p < 0.05$.

Results

Liver inflammation criteria

The liver IL-6 concentration of the LPS group was significantly higher than that of the control group ($p < 0.001$). It was also identified all pretreatment doses of silymarin having a

protective effect against increasing of IL-6 level due to LPS treatment reflected in a lower concentration of IL-6 in the liver tissues ($p < 0.01$ for 10 mg/kg silymarin and $p < 0.001$ for 20 mg/kg silymarin).

Oxidative damage criteria

The liver malondialdehyde concentration of the LPS group was significantly higher than that of the control group ($p < 0.001$). It was also shown that both doses including 10 and 20 mg/kg of silymarin have a protective effect against increasing lipid peroxidation due to LPS treatment reflected in a lower concentration of malondialdehyde in the liver tissues ($p < 0.001$; Fig 1). It was also observed and the catalase activity in the liver tissues of LPS group significantly increased more than that of the control group ($p < 0.001$). The findings also revealed that the highest dose of silymarin

increases the catalase activity in the liver tissues compared to the LPS ($p < 0.01$).

Liver biochemical criteria

The results of silymarin extract on liver function criteria of the LPS- treated rats: In LPS group, serum AST concentration of the rats was higher than that of the control ones ($p < 0.01$; Fig 2). Treatment of the animals by two doses of silymarin attenuated the serum concentration of AST ($p < 0.01$, $p < 0.001$). LPS also increased serum ALT compared to the control group ($p < 0.001$; Fig 3). All two dose of silymarin prevented increasing of ALT concentration due to LPS administration ($p < 0.001$). Changes in weight and high-density lipoprotein level in various groups of rat are shown in tables 1 and 2 respectively.




Fig 1. Effects of LPS and silymarin treatment on malondialdehyde level in the liver. Silymarin and LPS was administered intraperitoneally. Data showed as mean \pm SD, * $p < 0.001$ comparison with control and LPS group. MDA= Malondialdehyde; LPS= Lipopolysaccharide; S= Silymarin.




Fig 2. Effects of LPS and silymarin treatment on AST concentration of the rats in the blood serum. Silymarin and LPS was administered intraperitoneally. Data showed as mean±SD, * p<0.01 comparison with LPS group. AST= Aspartate aminotransferase; LPS= Lipopolysaccharide; S= Silymarin.




Fig 3. Effects of LPS and silymarin treatment on ALT concentration of the rats in the blood serum. Silymarin and LPS was administered intraperitoneally. Data showed as mean±SD, * p<0.001 comparison with control and LPS group. ALT= Alanine aminotransferase; LPS= Lipopolysaccharide; S= Silymarin.

Table 1. Changes in weight of various groups with each other

Tukey's multiple comparisons test	Summary
Control vs. silymarin 10 (mg/kg)	NS
Control vs. silymarin 20 (mg/kg)	NS
Silymarin 10 (mg/kg) vs. 2 (mg/kg) LPS	<0.01
Silymarin 20 (mg/kg) vs. 2 (mg/kg) LPS	<0.001
Control vs. 2 (mg/kg) LPS	NS

LPS= Lipopolysaccharide; NS= Not significant

Table 2. Changes in high-density lipoprotein level of silymarin groups with each other

Tukey's multiple comparisons test	Summary
Control vs. silymarin 10 (mg/kg)	NS
Control vs. silymarin 20 (mg/kg)	NS
Silymarin 10 (mg/kg) vs. 2 (mg/kg) LPS	<0.01
Silymarin 20 (mg/kg) vs. 2 (mg/kg) LPS	<0.001
Control vs. 2 (mg/kg) LPS	NS

LPS= Lipopolysaccharide; NS= Not significant

Western blot analysis

The results of western blot analysis are described as follows: protein expression of Bax/Bcl₂ was up-regulated in LPS group in the liver (p<0.001; Fig 4 and 5). Similarly, protein expressions of caspase-3, caspase-8 and caspase-9 were up-regulated by LPS in the liver (p<0.001; Fig 6-8). The data show significant results in silymarine 10 and 20 mg/kg group

and LPS group on the protein levels of Bax in the rat liver tissue. Representative western blots showed specific bands for Bcl2 and β-actin as an internal control. Representative western blots showed specific bands for Caspase-8 and Caspase-9 with β-actin as an internal control. Western blot analysis was conducted on the basis of other studies [27].




Fig. 4A. Effect of silymarin 10 and 20 mg/kg and LPS on the protein levels of Bax in the rat liver tissue. 1: saline, 2: LPS, 3: 10 mg/kg silymarin before 2 mg/kg LPS, 4: received 20 mg/kg silymarin before 2 mg/kg LPS, 5: LPS.




Fig. 4B. Effect of silymarin 10 and 20 mg/kg and LPS on the protein levels of Bax in the rat liver tissue. Densitometry data of protein analysis. Data are expressed as the mean±SD. *p<0.001 2 doses of silymarin in comparison with control and LPS group. LPS= Lipopolysaccharide; S= Silymarin.




Fig. 5A. Effect of silymarin 10 and 20 mg/kg and LPS on the protein levels of Bcl2 in the rat liver tissue. 1: saline, 2: 10 mg/kg silymarin before 2 mg/kg LPS, 3: received 20 mg/kg silymarin before 2 mg/kg LPS, 4: LPS, 5: LPS.




Fig. 5B. Effect of silymarin 10 and 20mg/kg and LPS on the protein levels of Bcl2 in the rat liver tissue. Densitometry data of protein analysis. Data are expressed as the mean±SD. *p<0.001 comparison with LPS group, **p<0.001 comparison with control group. LPS= Lipopolysaccharide; S= Silymarin.




Fig. 6A. Effect of silymarin 10 and 20 mg/kg and LPS on the protein levels of Caspase-3 in the rat liver tissue 1: saline, 2: LPS, 3: LPS, 4: 10 mg/kg silymarin before 2 mg/kg LPS, 5: received 20 mg/kg silymarin before 2 mg/kg LPS.




Fig. 6B. Effect of silymarin 10 and 20 mg/kg and LPS on the protein levels of Caspase-3 in the rat liver tissue. Densitometry data of protein analysis. Data are expressed as the mean±SD. *p<0.001 comparison with LPS group, **p<0.001 comparison with control and LPS group. LPS= Lipopolysaccharide; S= Silymarin.



Fig. 7A. Effect of silymarin 10 and 20 mg/kg and LPS on the protein levels of Cleaved-Caspase-8 in the rat liver tissue. 1: saline, 2: LPS, 3: LPS, 4: 10 mg/kg S before 2 mg/kg LPS, 5: received 20 mg/kg silymarin before 2 mg/kg LPS.




Fig. 7B. Effect of silymarin 10 and 20 mg/kg and LPS on the protein levels of Cleaved-Caspase-8 in the rat liver tissue. Densitometry data of protein analysis. Data are expressed as the mean±SD. *p<0.001 comparison with control group, **p<0.001 comparison with LPS group. LPS= Lipopolysaccharide; S= Silymarin.




Fig. 8A. Effect of silymarin 10 and 20 mg/kg and LPS on the protein levels of Caspase-9 in the rat liver tissue. 1: saline, 2: LPS, 3: LPS, 4: 10 mg/kg silymarin before 2 mg/kg LPS, 5: received 20 mg/kg silymarin before 2 mg/kg LPS.




Fig. 8B. Effect of silymarin 10 and 20 mg/kg and LPS on the protein levels of Caspase-9 in the rat liver tissue. Densitometry data of protein analysis. Data are expressed as the mean±SD. *p<0.001 comparison with LPS and control group, **p<0.001 comparison with LPS and control group. LPS= Lipopolysaccharide; S= Silymarin.

Discussion

Acute inflammation results in a general reaction of the organism which is followed by anorexia, loss of body weight, hypoglycemia, and changes in the levels of several plasma proteins produced by the hepatocytes. These conditions can also be reproduced by a number of experimental treatments. In particular, intra-peritoneal injection of LPS is commonly used to reproduce systemic inflammation [28, 29]. Administration of LPS for the animals is used as an experimental model to analyze the mechanism(s) underlying endotoxin-induced acute liver injury since it induces the infiltration of inflammatory cells into the liver and causes acute liver injury. In this study, we showed that LPS induces liver inflammation presented by an increased level of IL-6 in the liver tissues. LPS is also known to enhance the formation of ROS and lipid peroxidation products such as superoxide anions and peroxides and their secondary product, malondialdehyde, in liver [30]. The present data provided evidence that LPS injection increases malondialdehyde and decreases catalase activity. In another study, LPS promoted hepatic oxidative stress in rats. Livers of rats exposed to LPS exhibited elevated levels of lipid peroxidation products, malondialdehyde and hydro peroxides [31]. It has also been reported that the mice lacking iNOS gene are protected against acetaminophen-caused liver injury. Considering these evidences, an overproduction of NO may have a role in hepatic injury as observed in the recent study [32].

The measurement of hepatic enzymes appearing in the blood has been employed as a reliable indicator for assessment of hepatotoxicity. Our data revealed that the liver enzymes including ALT and AST are increased subsequent to inflammation induced by LPS. Consistently, LPS injection was previously reported that was followed by increased levels of AST and ALT, as well as apoptotic and necrotic changes in hepatocytes, which are biochemical and histological parameters of liver damage, respectively. Cytokines have also an important function in the development of hepatic inflammation and fibrosis [33, 34]. The results of the current study exhibited that silymarin improves liver functions in LPS injected rats presented by decreased levels of ALT, AST in the serum. Similar to the results, it has also been previously reported that silymarin ameliorates the development of ALT, AST levels in hepatic injury models [35, 36]. It has been demonstrated that silymarin inhibits LPS-induced production of pro-inflammatory cytokines such as IL-1 β in the liver. To evaluate the mechanism of the protective effects of silymarin in hepatic inflammation, we measured the levels of IL-6 in supernatants of liver tissues from each study group [37]. The results projected that the both groups treated with silymarin possessed lower IL-6 level in liver tissue compared to LPS group. The results also explored that the highest dose was more effective than other dose. These findings confirm the previous reports on the antioxidant effect of silymarin active

ingredients and compounds (Silybum) [38]. In the current study, our data revealed that Silybum inhibites the LPS-induced production of Interleukin, the activation of malondialdehyde and the caspase3-8-9, thus protecting mice against LPS-induced inflammation [39].

In the present study, we measured protein levels of Bax/Bcl₂ ratio, cleaved caspase-3, cleaved caspase-8 and cleaved caspase-9 for evaluation of apoptosis. Bcl₂ is an anti-apoptotic protein localized in the outer mitochondrial membrane, nuclear membrane and endoplasmic reticulum. Caspases are proteolytic enzymes which trigger cell death by cleaving specific proteins in the cytoplasm and nucleus. The initiator caspase-8 and caspase-9 may be stimulated by cell membrane death receptors thus leading to activation of caspase-3 by extrinsic pathways of apoptosis [40].

References

- [1]. Bissell DM, Gores GJ, Laskin DL, Hoofnagle JH. Drug-induced liver injury: mechanisms and test systems. *Hepatology*. 2001;33(4):1009-1013.
- [2]. Kono H, Rusyn I, Yin M, Gäbele E, Yamashina S, Dikalova A, et al. NADPH oxidase-derived free radicals are key oxidants in alcohol-induced liver disease. *J Clin Invest*. 2000; 106(7): 867-72.
- [3]. Shimada T, Yamazaki H, Mimura M, Inui Y, Guengerich FP. Interindividual variations in human liver cytochrome P-450 enzymes involved in the oxidation of drugs, carcinogens and toxic chemicals: studies with liver microsomes of 30 Japanese and 30 Caucasians. *J Pharmacol Experiment Therapeut*. 1994; 270(1): 414-23.
- [4]. Miwa M, Ura M, Nishida M, Sawada N, Ishikawa T, Mori K, et al. Design of a novel oral fluoropyrimidine carbamate, capecitabine, which generates 5-fluorouracil selectively in tumours by enzymes concentrated in human liver and cancer tissue. *Euro J Cancer* 1998; 34(8): 1274-281.
- [5]. Tilg H, Moschen AR. Evolution of inflammation in nonalcoholic fatty liver disease: the multiple parallel hits hypothesis. *Hepatology*. 2010; 52(5): 1836-846.
- [6]. Sewerynek E, Reiter RJ, Melchiorri D, Ortiz GG, Lewinski A. Oxidative damage in the liver induced by ischemia-reperfusion: protection by melatonin. *Hepato-gastroenterol*. 1996; 43(10): 898-905.
- [7]. Beumer C, Wulferink M, Raaben W, Fiechter D, Brands R, Seinen W. Calf intestinal alkaline phosphatase, a novel therapeutic drug for lipopolysaccharide (LPS)-mediated diseases, attenuates LPS toxicity in mice and piglets. *J Pharmacol Experiment Therapeut*. 2003; 307(2): 737-44.
- [8]. Lehmann V, Freudenberg M, Galanos C. Lethal toxicity of lipopolysaccharide and tumor necrosis

Conclusion

The results of this research demonstrated that silymarin can exert protective effects against toxic effects of LPS in rat liver such as IL-6 and malondialdehyde. Anti-inflammatory drug can play a protective role in attenuating the liver inflammation induced by LPS injection. LPS also increased expression of Bax/Bcl₂ and caspases ratio as apoptosis parameters. This study suggests silymarin as the most important liver protective compound.

Conflict of Interest

No potential conflict of interest was reported by the authors.

Acknowledgments

The writers of this study would like to express their gratitude to all the hardworking staff of the Microbiology and Biotechnology Research Center for their valuable support. This research was financially sponsored by Shahrekord Islamic Azad University in Shahrekord, Iran (2018-239). The study was financially supported by both the personal as well as Islamic Azad University of Shahrekord, Iran.

- factor in normal and D-galactosamine-treated mice. *J Experiment Med.* 1987; 165(3): 657-63.
- [9]. Xu DX, Chen YH, Zhao L, Wang H, Wei W. Reactive oxygen species are involved in lipopolysaccharide-induced intrauterine growth restriction and skeletal development retardation in mice. *Am J Obstetrics Gynecol.* 2006; 195(6): 1707-714.
- [10]. Koyama I, Matsunaga T, Harada T, Hokari S, Komoda T. Alkaline phosphatases reduce toxicity of lipopolysaccharides in vivo and in vitro through dephosphorylation. *Clin Biochem.* 2002; 35(6): 455-61.
- [11]. Li J, Baud O, Vartanian T, Volpe JJ, Rosenberg PA. Peroxynitrite generated by inducible nitric oxide synthase and NADPH oxidase mediates microglial toxicity to oligodendrocytes. *Proceedings of the National Acad Sci.* 2005; 102(28): 9936-941.
- [12]. Azizi-Malekabadi H, Abareshi A, Beheshti F, Marefati N, Norouzi F, Soukhtanloo M, et al. The effect of captopril on inflammation-induced liver injury in male rats. *Toxin Rev.* 2018; 1-9.
- [13]. Shaker E, Mahmoud H, Mnaa S. Silymarin, the antioxidant component and *Silybum marianum* extracts prevent liver damage. *Food and Chem Toxicol.* 2010; 48(3): 803-806.
- [14]. Bosisio E, Benelli C, Pirola O. Effect of the flavanolignans of *Silybum marianum* L. on lipid peroxidation in rat liver microsomes and freshly isolated hepatocytes. *Pharmacol Res.* 1992; 25(2): 147-65.
- [15]. Patel N, Joseph C, Corcoran GB, Ray SD. Silymarin modulates doxorubicin-induced oxidative stress, Bcl-xL and p53 expression while preventing apoptotic and necrotic cell death in the liver. *Toxicol applied pharmacol.* 2010; 245(2): 143-52.
- [16]. Anaeigoudari A, Shafei MN, Soukhtanloo M, Sadeghnia HR, Reisi P, Beheshti F, et al. Lipopolysaccharide-induced memory impairment in rats is preventable using 7-nitroindazole. *Arquivos de Neuro-psiquiatria.* 2015; 73(9): 784-90.
- [17]. Hosseini M, Pourganji M, Khodabandehloo F, Soukhtanloo M, Zabihi H. Protective effect of L-arginine against oxidative damage as a possible mechanism of its beneficial properties on spatial learning in ovariectomized rats. *Basic Clin Neurosci.* 2012; 3(5): 36-44.
- [18]. Yilmaz HR, Sogut S, Ozyurt B, Ozugurlu F, Sahin S, Isik B, et al. The activities of liver adenosine deaminase, xanthine oxidase, catalase, superoxide dismutase enzymes and the levels of malondialdehyde and nitric oxide after cisplatin toxicity in rats: protective effect of caffeic acid phenethyl ester. *Toxicol Industr Health.* 2005; 21(1-2): 67-73.
- [19]. Videla LA, Rodrigo R, Orellana M, Fernandez V, Tapia G, Quinones L, et al. Oxidative stress-related parameters in the liver of non-alcoholic fatty liver disease patients. *Clin Sci.* 2004; 106(3): 261-68.
- [20]. Akhgari M, Abdollahi M, Kebryaezadeh A, Hosseini R, Sabzevari O. Biochemical evidence for free radical-induced lipid peroxidation as a mechanism for subchronic toxicity of malathion in blood and liver of rats. *Human Experiment Toxicol.* 2003; 22(4): 205-11.
- [21]. Nyblom H, Berggren U, Balldin J, Olsson R. High AST/ALT ratio may indicate advanced alcoholic liver disease rather than heavy drinking. *Alcohol and alcoholism.* 2004; 39(4): 336-39.
- [22]. Lum G, Gambino SR. Serum gamma-glutamyl transpeptidase activity as an indicator of disease of liver, pancreas, or bone. *Clin Chem.* 1972; 18(4): 358-62.
- [23]. Sakinah SS, Handayani ST, Hawariah LA. Zerumbone induced apoptosis in liver cancer cells via modulation of Bax/Bcl-2 ratio. *Cancer Cell Int.* 2007; 7(1): 4.
- [24]. Panasiuk A, Dzieciol J, Panasiuk B, Prokopowicz D. Expression of p53, Bax and Bcl-2 proteins in hepatocytes in non-alcoholic fatty liver disease. *World J Gastroenterol.* 2006; 12(38): 6198.
- [25]. Kren BT, Trembley JH, Krajewski S, Behrens TW, Reed JC, Steer CJ. Modulation of apoptosis-associated genes bcl-2, bcl-x, and bax during rat liver regeneration. *Cell Growth Differentiation-Publica Am Associ Cancer Res.* 1996; 7(12): 1633-642.
- [26]. Tsujimoto Y, Shimizu S. Bcl-2 family: life-or-death switch. *FEBS Lett.* 2000; 466(1): 6-10.
- [27]. Boussabbeh M, Salem IB, Belguesmi F, Neffati F, Najjar MF, Abid-Essefi S, et al. Crocin protects the liver and kidney from patulin-induced apoptosis in vivo. *Environment Sci Pollution Res.* 2016; 23(10): 9799-808.
- [28]. Nolan JP. Endotoxin, reticuloendothelial function, and liver injury. *Hepatol.* 1981; 1(5): 458-65.
- [29]. Simeonova PP, Gallucci RM, Hulderman T, Wilson R, Kommineni C, Rao M, et al. The role of tumor necrosis factor- α in liver toxicity, inflammation, and fibrosis induced by carbon tetrachloride. *Toxicol appl pharmacol.* 2001; 177(2): 112-20.
- [30]. Pulskamp K, Diabaté S, Krug HF. Carbon nanotubes show no sign of acute toxicity but induce intracellular reactive oxygen species in dependence on contaminants. *Toxicol Lett.* 2007; 168(1): 58-74.
- [31]. Lu Y, Zhuge J, Wang X, Bai J, Cederbaum AI. Cytochrome P450 2E1 contributes to ethanol-induced fatty liver in mice. *Hepatol.* 2008; 47(5): 1483-494.
- [32]. Bourdi M, Masubuchi Y, Reilly TP, Amouzadeh HR, Martin JL, George JW, et al.

- Protection against acetaminophen-induced liver injury and lethality by interleukin 10: role of inducible nitric oxide synthase. *Hepatology*. 2002; 35(2): 289-98.
- [33]. Zhang S, Yang N, Ni S, Li W, Xu L, Dong P, et al. Pretreatment of lipopolysaccharide (LPS) ameliorates D-GalN/LPS induced acute liver failure through TLR4 signaling pathway. *Int J Clin Experiment Pathol*. 2014; 7(10): 6626.
- [34]. Giriwono PE, Shirakawa H, Ohsaki Y, Hata S, Kuriyama H, Sato S, et al. Dietary supplementation with geranylgeraniol suppresses lipopolysaccharide-induced inflammation via inhibition of nuclear factor- κ B activation in rats. *Eur J Nutr*. 2013; 52(3): 1191-199.
- [35]. Lin MC, Kao SH, Chung PJ, Chan KC, Yang MY, Wang CJ. Improvement for high fat diet-induced hepatic injuries and oxidative stress by flavonoid-enriched extract from *Nelumbo nucifera* leaf. *J Agriculture Food Chem*. 2009; 57(13): 5925-932.
- [36]. Ball KR, Kowdley KV. A review of *Silybum marianum* (milk thistle) as a treatment for alcoholic liver disease. *Journal of Clinical Gastroenterology*. 2005; 39(6): 520-28.
- [37]. Naugler WE, Sakurai T, Kim S, Maeda S, Kim K, Elsharkawy AM, et al. Gender disparity in liver cancer due to sex differences in MyD88-dependent IL-6 production. *Science*. 2007; 317(5834): 121-24.
- [38]. Kovalovich K, Li W, DeAngelis R, Greenbaum LE, Ciliberto G, Taub R. Interleukin-6 protects against Fas-mediated death by establishing a critical level of anti-apoptotic hepatic proteins FLIP, Bcl-2, and Bcl-xL. *J of Biological Chemistry*. 2001; 276(28): 26605-6613.
- [39]. Barrier A, Olaya N, Chiappini F, Roser F, Scatton O, Artus C, et al. Ischemic preconditioning modulates the expression of several genes, leading to the overproduction of IL-1Ra, iNOS, and Bcl-2 in a human model of liver ischemia-reperfusion. *FASEB J*. 2005; 19(12): 1617-626.
- [40]. Sidi S, Sanda T, Kennedy RD, Hagen AT, Jette CA, Hoffmans R, et al. Chk1 suppresses a caspase-2 apoptotic response to DNA damage that bypasses p53, Bcl-2, and caspase-3. *Cell*. 2008; 133(5): 864-77.