

Original Article

Significant Impact of let-7d MicroRNA on Breast Cancer Cell Lines Post-Radiation Treatment

Mahdieh Nejadtaghi ¹ Ph.D., Hossein Mozdarani ^{1*} Ph.D.¹ Department of Medical Genetics, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

A B S T R A C T

Article history

Received: 31 Oct 2023

Accepted: 23 Jun 2024

Available online: 23 Sep 2024

Keywords

Breast Cancer

DNA Repair

MiRNA

Let-7d

Radio Resistance

Radiosensitivity

Introduction: Radiotherapy is a common treatment for breast cancer treatment, that induces DNA damage. These DNA damages are addressed through various repair pathways, which regulate the DNA repair systems and confer radio-resistance. Non-coding RNAs are a big proportion of genome transcripts without the potential to encode proteins. Related studies demonstrated that radiation affects the expression of non-coding RNAs. Let-7d, a tumor suppressor in numerous cancers, has some target genes that play a role in the DNA repair system.

Materials and Methods: Human breast cancer cell lines MDA-MB-231 and MCF-7 were cultured in a Dulbecco's Modified Eagle Medium. The exponentially growing cells were treated with some doses of X-rays. After radiation treatment and cell harvesting, RNA was extracted, and cDNA synthesis was done. The let-7d miRNA expression changes were calculated with real-time quantitative reverse transcription polymerase chain reaction.

Results: The results implied that radiation caused increased let-7d expression in breast cancer cell lines after radiation treatment. In addition, the results showed that 24 h after radiation, the expression of let-7d in the radioresistant cell line was higher than the radiosensitive one; 48 h after radiation, the expression of let-7d in the radiosensitive cell line was higher than the other one.

Conclusions: The results demonstrated that radiation treatment increased let-7d miRNA expression in both radiosensitive and radio-resistant breast cancer cell lines. Therefore, let-7d might be involved in the radiosensitivity of breast cancer.



CC BY-NC 4.0

* **Corresponding Author:** Department of Medical Genetics, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran. **Tel:** +989121374831, **Email:** mozdarah@modares.ac.ir

Introduction

Breast cancer is the second leading cause of cancer deaths and the most common malignancy in women. According to statistics, there is a significant rise in breast cancer incidence, especially in Iran, and the average age of onset in Iranian women is 45 years, while this figure is at least ten years higher in other countries [1, 2]. At present, radiotherapy is the main means for patients with advanced breast cancer [3]. Ionizing radiation has a destroying effect in the cells called Double Strand Breaks (DSBs), so the DNA repair pathways are involved in response to radiotherapy [4]. The most important DNA repair pathways for repairing DNA strand breaks are homologous recombination and non-homologous end joining (NHEJ) [5-7]. Non-coding RNAs constitute a large fraction of genome transcripts without being able to code for proteins. Which can be divided into small (200 nt) and long ncRNAs (lncRNAs) [8, 9]. lncRNAs range from 200 nucleotides to less than 100 kb in length [10, 11]. miRNAs are a class of small non-coding RNA molecules that negatively regulate gene expression at the post-transcriptional level by altering the stability or translational efficiency of its target mRNAs [12, 13]. In the context of the DNA damage response, miRNAs may determine cell fate by dictating the appropriate cellular outcome in response to different levels of DNA damage [14]. In support of this, recent studies have shown expression patterns of miRNAs that are associated with increasing

doses of ionizing radiation and ultraviolet [15, 16].

The let-7 family of miRNAs play a vital role in regulating carcinogenesis; one of its members is let-7d, which is located on chromosome 9q22.3, which acts as a tumor suppressor in numerous human cancers [17]. This miRNA has some target genes that play a role in the DNA repair system [18-20]. One study demonstrated that in radioresistant and radiosensitive glioblastoma cell lines, the let-7d expression was higher in radioresistant cell line. Also, they showed that upregulation of let-7d caused radioresistance [21]. In another study, the results suggested the role of let-7d in the radiosensitivity of breast cancer stem cells [22]. Evaluating let-7d expression changes in response to radiation in endothelial cells showed its upregulation after receiving radiation [23]. Also, let-7d plays a role in the sensitivity of hypopharyngeal carcinoma cells to radiation and chemotherapeutic drugs [24]. Also, its shown that let-7d could suppress the Epithelial-mesenchymal transition pathway [25], and its demonstrated that this pathway could cause radioresistance in many cancers [26].

The goal of the present study was to evaluate let-7d miRNA expression changes under the effect of different radiation doses in MCF-7 and MDA-MB- 231 breast cancer cell lines that have somewhat inherent sensitivity and resistance to radiation respectively, to explore let-7d potential as novel markers for irradiation in breast cancer.

Materials and Methods

Cell Culture

MCF7 and MDA-MB- 231 breast cancer cell lines were obtained from the Pasteur Institute of Iran (Tehran, Iran) and cultured at 37 °C in humidified 5% CO₂. All the cell lines were cultured in Dulbecco's Modified Eagle (DMEM Gibco) medium supplemented with 10% fetal bovine serum (FBS) (Bio-Idea, Iran) and 1% penicillin/ streptomycin (PAN-Biotech GmbH).

Irradiation

Cells were irradiated using a 6-MeV linear accelerator (LINAC) (Elekta, Sweden) with 2, 4, and 8 Gy radiation doses, and we had a control flask without receiving any radiation dose. For all the radiation treatments and the control sample, we had two flasks; one group of flasks was incubated for the next 24 h, and one group was incubated for the next 48 h. The radiation doses were selected based on related articles. Finally, after trypsinizing, the cells were harvested. We performed the experiments in triplicate.

RNA isolation and quality control

Total RNA was extracted from cells using TRIzol reagent (Geneall, South Korea) according to the manufacturer's instructions. The concentration and quality of the isolated RNA were assessed on a Nanodrop ND-1000 spectrophotometer. All extracted RNAs were stored at -80°C.

Real-time polymerase chain reaction (RT-PCR) and primer designing

For cDNAs synthesis, a reverse transcription kit (Beta Bayern, Germany) was used according to the manufacturer's instructions. Stem-loop primers were used for the reverse transcription of let-7d miRNA.

The RT-PCR was performed using SYBR green with Applied Biosystems instrument in duplicate by adopting the $2^{-\Delta\Delta Ct}$ method. The expression levels of the miRNA were normalized to the endogenous control U6. The specific primer sequences are shown in table 1.

Statistical Analysis

Statistical analyses were performed using GraphPad Prism 8 software. The data were expressed as means \pm SD. The one-way ANOVA test evaluated the difference between groups. For comparing MCF-7 and MDA-MB-231 cell lines, the independent t-test was used for each dose of radiation.

Results

To study the effects of irradiation on let-7d miRNA expression in MCF-7 and MDA-MB-231 cells, we employed Real-time PCR before and after receiving different radiation doses. Finally, the results suggested up-regulation of let-7d miRNA expression after irradiation.

Table 1. The sequence of primers

Gene	Forward primer	Reverse primer
Let-7d	ACGCAGAGGTAGTAGGTTGC	TGCAGGGTCCGAGGTATTTCG
U6	CTCGCTTCGGCAGCACATATACT	ACGCTTCACGAATTTGCGTGTC

Let-7d Expression changes in response to radiation in the MCF-7 cell line

let-7d expression was up-regulated in the MCF-7 cell line at 24 and 48 hours post-irradiation compared to the control sample. Approximately 24 hours after irradiation, the clinically relevant dose of 8 Gy ($p < 0.01$) induced significant changes in let-7d miRNA levels. 48 h after irradiation, there was up-regulation in 2 ($p < 0.001$), 4 ($p < 0.001$) and 8 Gy ($p < 0.0001$) that peaked at 8 Gy (Figure 1 A, B).

Let-7d expression changes in response to radiation in the MDA-MB-231 cell line

As observed in the MCF-7 cell line, let-7d expression was up-regulated in the MDA-MB-231 cell line 24 h and 48h after irradiation in comparison with the control sample. 24 h after irradiation, clinically relevant doses of 2 ($p < 0.001$), 4 ($p < 0.0001$) and 8 Gy ($p < 0.001$) induced significant changes in let-7d miRNA levels. This upregulation peaked at 4 Gy. 48 hours after irradiation, there was up-regulation in 2 ($p < 0.001$) and 4 ($p < 0.05$) Gy that peaked at 2 Gy (Fig.1 C, D).




Fig. 1. The let-7d miRNA expression changes after receiving different radiation doses in the MCF-7 cell line after 24 h (A) and 48 h (B). And in the MDA-MB-231 cell line after 24 h (C) and 48 h (D). * $p < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

A comparison between the relative expression of let-7d miRNA in MCF-7 and MDA-MB-231 cell lines

The relative expression of let-7d miRNA 24 h after receiving irradiation in the MDA-MB-231 cell line compared with the MCF-7 cell line was significantly higher for all doses of radiation. This difference was

more pronounced at 4 Gy. These findings are demonstrated in Figure 2.

However, 48 h after receiving irradiation the let-7d expression in the MCF-7 cell line was significantly higher than the MDA-MB-231 cell line. At this time, the difference was more pronounced at 8 Gy.




Fig. 2. A Comparison of the relative expressions of let-7d miRNA between MCF-7 and MDA-MB-231 cell lines in each radiation dose 24 h and 48 h after receiving radiation. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $P < 0.0001$.

Discussion

breast cancer is among the most frequent causes of women's death worldwide. Radiotherapy is one of the mainstream approaches for its treatment. Radiation promotes a series of changes in tumor cells. DNA damage response is one of the main factors which regulate cell survival after radiation [27]. So, insight into the effect of radiation on DNA damage pathways provides an improvement in the prognosis of cancer patients [28].

In recent years, non-coding RNAs, especially miRNAs, as a hot spot, have attracted widespread concern [29]. MicroRNAs are a class of regulatory molecules, and miRNAs have an emerging role in modulating cell physiology and different pathways like DNA repair. It described that miRNAs could be induced by radiation, playing a role as intermediate molecules in the regulation of DNA damage response [30, 31].

Several previous studies have indicated that there is a potential for plasma miRNAs as valuable biomarkers of radiation exposure. A growing body of evidence indicates that let-7d miRNA has expression changes in response to radiation and plays a pivotal role in radio sensitivity or radioresistance of tumor cells.

In order to show let-7d role in the etiology of breast cancer, one study by Wei et al. showed that let-7d was downregulated in breast cancer tissues, and also, they showed that let-7d inhibits growth and metastasis in breast cancer [32].

In the present study, we determined the expression levels of let-7d miRNA in MCF-7 and MDA-MB231 breast cancer cell lines under the effect of radiation and found that let-7d was

overexpressed in the irradiated group as compared with controls. Some other studies are showing similar changes in let-7d expression produced on radiated cells. For example, in the study by Lamperska et al., increased levels of this miRNA expression were shown under the effect of radiation in irradiated hypopharynx cancerous cells [24]. In another study, the results suggested an upregulation of let-7d expression after radiation to endothelial cells [23].

To evaluate the relationship between let-7d and radiosensitivity/resistance of breast cancer, a comparison between the expression level between MCF-7 and MDA-MB-231 cell lines was performed. The higher expression level of let-7d in the radioresistant cell line 24 h after receiving radiation and the higher expression level in the radiosensitive cell line 48 h after receiving radiation suggested that besides playing a role in radiosensitivity, let-7d miRNA would contribute to the radioresistance of breast cancer. However, the mechanism of how this miRNA plays these roles is not clear yet. Basic research on its function on radiosensitivity/resistance of breast cancer is urgently needed. However, probably because of the role of let-7d in both the DNA repair pathway and apoptosis [33], 24 h after receiving radiation, its function in the DNA repair pathway is more activated. Hence, its expression level was higher in radioresistance cell line and 48 h after radiation because of the severity of damage its role in apoptosis is more activated, so the expression level in a radiosensitive cell line is higher than the other one. Different studies are showing the

role of let-7d in the radiosensitivity/ resistance of various cancers. For example, one study demonstrated that in glioblastoma, its upregulation caused radioresistance [21]. Alternatively, one study suggested the role of let-7d in the radiosensitivity of breast cancer stem cells [22].

Conclusion

This study's results suggested that let-7d miRNA has the potential to be a radiation marker in breast cancer; in addition, maybe it could play a role in the radiosensitivity/resistance of breast cancer patients. However, verifying these findings needs functional studies, in addition to evaluating this miRNA expression in breast cancer patients.

References

- [1]. Vostakolaei FA, Broeders MJ, Rostami N, Van Dijck JA, Feuth T, Kiemeny LA, et al. Age at diagnosis and breast cancer survival in Iran. *International Journal of Breast Cancer* 2012; 2012: 517976.
- [2]. Farhood B, Raei B, Malekzadeh R, Shirvani M, Najafi M, Mortezaadeh T. A review of incidence and mortality of colorectal, lung, liver, thyroid, and bladder cancers in Iran and compared to other countries. *Contemporary Oncology* 2019; 23(1): 7-15.
- [3]. Gupta N, Chugh Y, Chauhan AS, Pramesh C, Prinja S. Cost-effectiveness of Post-Mastectomy Radiotherapy (PMRT) for breast cancer in India: An economic modelling study. *The Lancet Regional Health-Southeast Asia*. 2022; 4(1): 1-7.
- [4]. Zhao L, Bao C, Shang Y, He X, Ma C, Lei X, et al. The determinant of DNA repair pathway choices in ionising radiation-induced DNA double-strand breaks. *BioMed Research International* 2020; 2020: 4834965.
- [5]. Zhu J, Chen S, Yang B, Mao W, Yang X, Cai J. Molecular mechanisms of lncRNAs in regulating cancer cell radiosensitivity. *Bioscience Reports* 2019; 39(8): 590.
- [6]. Murmann-Konda T, Soni A, Stuschke M, Iliakis G. Analysis of chromatid-break-repair detects a homologous recombination to non-homologous end-joining switch with increasing load of DNA double-strand breaks. *Mutation Research/ Genetic Toxicology and Environmental Mutagenesis* 2021; 867: 503372.
- [7]. Nickoloff JA, Sharma N, Allen CP, Taylor L, Allen SJ, Jaiswal AS, et al. Roles of homologous recombination in response to ionizing radiation-induced DNA damage. *International Journal of Radiation Biology* 2023; 99(6): 903-14.
- [8]. Xu D, Yuan W, Fan C, Liu B, Lu MZ, Zhang J. Opportunities and challenges of predictive approaches for the non-coding RNA in Plants. *Frontiers in Plant Science* 2022; 414: 890663.
- [9]. He J, Wang HB, Huang JJ, Zhang L, Li DL, He WY, et al. Diabetic neuropathic pain induced by streptozotocin alters the expression profile of non-coding RNAs in the spinal cord of mice as determined by sequencing analysis. *Experimental and Therapeutic Medicine* 2021; 22(1): 1-11.
- [10]. Dhanoa JK, Sethi RS, Verma R, Arora JS, Mukhopadhyay CS. Long non-coding RNA: its evolutionary relics and biological implications in

Ethical Considerations

All ethical considerations were followed in compiling this work. This study was supported by the Research Department of Tarbiat Modares University (grant number: IG-39711, ethics code: (IR.MODARES.REC.1399.010), Tehran, Iran.

Funding

This research has been supported by Tarbiat Modares University Tehran, Iran, by grant number IG-39711 provided to Prof. H. Mozdarani.

Conflict of Interest

The authors declare that there is no conflict of interest regarding this article's publication and financial issues.

Acknowledgment

The authors also thank Mr. Hassan Nosrati for the irradiation of samples.

Authors' Contributions

MN performed experiments, sample collection, data curation, and data analysis and wrote the manuscript draft. HM designed and supervised the research plan, prepared, analyzed data analysis, edited and approved the final manuscript.

- mammals: a review. *Journal of Animal Science and Technology* 2018; 60: 1-10.
- [11]. Li J, Xuan Z, Liu C. Long non-coding RNAs and complex human diseases. *International Journal of Molecular Sciences* 2013; 14(9): 18790-8808.
- [12]. El-Serag HB. Hepatocellular carcinoma: an epidemiologic view. *Journal of Clinical Gastroenterology* 2002; 35(S5): 72-8.
- [13]. Valinezhad Orang A, Safaralizadeh R, Kazemzadeh-Bavili M. Mechanisms of miRNA-mediated gene regulation from common downregulation to mRNA-specific upregulation. *Int J Genomics*. 2014; 2014: 970607.
- [14]. Visser H, Thomas AD. MicroRNAs and the DNA damage response: How is cell fate determined? *DNA Repair*. 2021; 108: 103245.
- [15]. Liao XH, Zheng L, He HP, Zheng DL, Wei ZQ, Wang N, et al. STAT3 regulated ATR via microRNA-383 to control DNA damage to affect apoptosis in A431 cells. *Cellular Signalling* 2015; 27(11): 2285-295.
- [16]. Harding SM, Coackley C, Bristow RG. ATM-dependent phosphorylation of 53BP1 in response to genomic stress in oxic and hypoxic cells. *Radiotherapy and Oncology* 2011; 99(3): 307-12.
- [17]. Wang Z, Lin S, Li JJ, Xu Z, Yao H, Zhu X, et al. MYC protein inhibits transcription of the microRNA cluster MC-let-7a-1~ let-7d via noncanonical E-box. *Journal of Biological Chemistry* 2011; 286(46): 39703-9714.
- [18]. Büssing I, Slack FJ, Großhans H. let-7 microRNAs in development, stem cells and cancer. *Trends in Molecular Medicine* 2008; 14(9): 400-409.
- [19]. Luoto KR, Meng AX, Wasylishen AR, Zhao H, Coackley CL, Penn LZ, et al. Tumor cell kill by c-MYC depletion: role of MYC-regulated genes that control DNA double-strand break repair. *Cancer Research* 2010; 70(21): 8748-759.
- [20]. Li AY, Boo LM, Wang SY, Lin HH, Wang CC, Yen Y, et al. Suppression of non-homologous end joining repair by overexpression of HMGA2. *Cancer Research* 2009; 69(14): 5699-706.
- [21]. Matos B, Boštjančič E, Popović M, Glavač D. Dynamic of expression of 11 microRNAs in recurrent glioblastoma before and after treatment with radiotherapy and chemotherapy. *Journal of Neurological Surgery Part A: Central European Neurosurgery* 2015; 76(S 02): 100.
- [22]. Sun H, Ding C, Zhang H, Gao J. Let - 7 miRNAs sensitize breast cancer stem cells to radiation-induced repression through inhibition of the cyclin D1/Akt1/Wnt1 signaling pathway. *Molecular Medicine Reports* 2016; 14(4): 3285-292.
- [23]. Wagner-Ecker M, Schwager C, Wirkner U, Abdollahi A, Huber PE. MicroRNA expression after ionizing radiation in human endothelial cells. *Radiation Oncology* 2010; 5(1): 1-10.
- [24]. Lamperska KM, Kolenda T, Teresiak A, Kowalik A, Kruszyna-Mochalska M, Jackowiak W, et al. Different levels of let-7d expression modulate response of FaDu cells to irradiation and chemotherapeutics. *PLoS One* 2017; 12(6): 180265.
- [25]. Chang CJ, Hsu CC, Chang CH, Tsai LL, Chang YC, Lu SW, et al. Let-7d functions as novel regulator of epithelial-mesenchymal transition and chemoresistant property in oral cancer. *Oncology Reports* 2011; 26(4): 1003-1010.
- [26]. Tan J, Qiu K, Li M, Liang Y. Double-negative feedback loop between long non-coding RNA TUG1 and miR-145 promotes epithelial to mesenchymal transition and radio-resistance in human bladder cancer cells. *FEBS Letters* 2015; 589(20): 3175-181.
- [27]. Wang Y, Wang Q, Chen S, Hu Y, Yu C, Liu R, et al. Screening of long non-coding RNAs induced by radiation using microarray. *Dose-Response* 2020; 18(2): 155.
- [28]. Wengner AM, Scholz A, Haendler B. Targeting DNA damage response in prostate and breast cancer. *International Journal of Molecular Sciences* 2020; 21(21): 8273.
- [29]. Lopez-Bertoni H, Latorra J. Opinion: miRNAs—The new wave of molecular cancer therapeutics. *Translational Oncology* 2021; 14(6): 1064.
- [30]. Wan G, Mathur R, Hu X, Zhang X, Lu X. miRNA response to DNA damage. *Trends in Biochemical Sciences* 2011; 36(9): 478-84.
- [31]. Han C, Wan G, Langley RR, Zhang X, Lu X. Crosstalk between the DNA damage response pathway and microRNAs. *Cellular and Molecular Life Sciences* 2012; 69: 2895-906.
- [32]. Wei Y, Liu G, Wu B, Yuan Y, Pan Y. Let-7d inhibits growth and metastasis in breast cancer by targeting Jab1/Cops5. *Cellular Physiology and Biochemistry* 2018; 47(5): 2126-135.
- [33]. Chen YN, Ren CC, Yang L, Nai MM, Xu YM, Zhang F, et al. MicroRNA let - 7d - 5p rescues ovarian cancer cell apoptosis and restores chemosensitivity by regulating the p53 signaling pathway via HMGA1. *International Journal of Oncology* 2019; 54(5): 1771-784.