

Original Article

Pulmonoprotective Action of *Zataria Multiflora* in an Experimental Model of Lung Inflammation Induced by Ochratoxin-A in Mice

Fahimeh Nourbakhsh¹ Ph.D., Hossein Chahardah Masoumi¹ Ph.D.,
Mohammadreza Saebi^{2*} Ph.D., Elaheh Tajbakhsh³ Ph.D.

¹ Deputy of Food and Drug, Isfahan University of Medical Sciences, Isfahan, Iran

² Department of Food Hygiene, Faculty of Veterinary Medicine, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran

³ Department of Microbiology, Faculty of Basic Sciences, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran

ABSTRACT

Article history

Received: 1 Oct 2024

Accepted: 9 Dec 2025

Available online: 22 Apr 2026

Keywords

Apoptosis

Apoptosis pathway

Lung inflammation

Ochratoxin A

Zataria multiflora



[CC BY-NC 4.0](https://creativecommons.org/licenses/by-nc/4.0/)

© Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License, which permits any non-commercial use, sharing, adaptation, distribution, and reproduction in any medium or format, provided that appropriate credit is given to the original author(s) and the source.

Introduction: Ochratoxin-A (OTA) represents one of the primary secondary metabolites produced by *Penicillium* and *Aspergillus* species. Acratoxin was recognized in 1965 as the most significant carcinogenic mycotoxin found in agricultural products. Given that the vital function of the lungs in maintaining other bodily organ system, it was examined the role of the *Zataria multiflora* extract in an experimental model of lung inflammation induced by Ochratoxin-A in mice.

Materials and Methods: This study was designed around a 14-day gavage administration of OTA in mice. Markers of oxidative stress and apoptotic factors (Bax/Bcl2, Caspase 3 Caspase 9) were also evaluated.

Results: OTA-induced toxicity provokes lung inflammation and apoptosis in a dose-dependent manner. The administration of *Zataria multiflora* (at dosages of 25 and 50 mg/kg) resulted in a significant alleviated of lung inflammation, oxidative damage, inflammatory cytokines, and levels of apoptosis protein when compared to the group treated with OTA alone.

Conclusion: The results of the study indicate that *Zataria multiflora* exerts robust antioxidant effects that counteract the inflammation and apoptosis associated with OTA-induced toxicity in lung tissue.

* **Corresponding Author:** Department of Food Hygiene, Faculty of Veterinary Medicine, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran. **Email:** msaebi0311@gmail.com

Introduction

Mycotoxins are among the most important metabolites produced by fungi and can exert diverse physiological effects [1]. Mycotoxins come in a variety of forms; six groups of them have significant toxic effects produced by *Fusarium*, *Alternaria*, and *Penicillium* and *Aspergillus* groups [2]. Ochratoxin-A (OTA) produced by a broad spectrum of fungi - especially *Aspergillus ochraceus*- is a white crystalline substance that dissolves readily in fats and polar organic solvents [3, 4]. OTA has been reported in a wide range of food sources, including cereals, coffee, dried fruits, grape juice, and meat products [5-7]. Nonetheless, cereals and cereal-based items are known the primary contributors to human exposure. OTA, which has the chemical formula C₂₀H₁₈O₆NCl, is regarded as a major mycotoxin that shows good solubility in polar organic solvents [8, 9]. According to the American Cancer Research Institute, OTA is categorized as a class B carcinogen. Among more than 300 known mycotoxins, OTA stands out as a key toxin with immunotoxic, and nephrotoxic properties [10, 11]. This toxin negatively affects the kidneys and liver and can cross the placenta; it also causes vasoconstriction and suppresses immune system activity [12, 13]. Due to its particular features related to cellular toxicity, OTA is considered as major contributor to infertility, including male infertility and fetal mortality [12, 14].

OTA inhibits cell proliferation, suppresses B and T lymphocytes, degrades immune organs, and decreases antibody production. It alters cytokine production, which affects both the number and

function of immune cells. OTA inhibits the death of immune cells by inhibiting protein synthesis by intensifying the death of immune cells and reducing their replacement rate [15].

Alongside its liver damage, nerve damage, immune damage, teratogenesis, and carcinogenesis of OTA, authors indicated that it easily effects on bronchial epithelial cells- within four hours of human exposure) [16, 17]. Various studies have shown that OTA specifically induces oxidative stress and activates the Nrf2 signaling pathway, ultimately leading to cytotoxicity and cell death [17, 18]. Another study by Okutan et al. similarly reported OTA's toxic effects on the lungs and heart, noting that oxidative stress and degradation of cellular proteins are the main mechanisms responsible for toxicity in these organs [19]. Despite numerous studies into OTA's toxicity, no study has yet clarified the molecular mechanisms OTA-induced lung damage or examined the protective effect of *Zataria multiflora* (*Z. multiflora*) In this context. *Z. multiflora* is a perennial plant known for its wide range of medical benefits and has attracted researchers' attention for many years [21-23]. Iran and Afghanistan are recognized as the primary regions for the growth and distribution of this species. Studies highlight their notable soothing and anti-inflammatory properties. In Iran, *Z. multiflora* is commonly used for its anti-inflammatory and antiseptic actions. The antioxidant effects of this plant are due to its most important active ingredient called Carvacrol [24, 25]. It affects the muscarinic

receptors and β -adrenoceptors receptors [26]. Numerous therapeutic effects have been demonstrated for both the plant and Carvacrol, especially regarding respiratory disorders. Another pharmacological action of Carvacrol is its ability to reduce cell death. Therefore, it was examined OTA-induced pulmonary inflammation and apoptosis in mice, as well as the protective effects of *Z. multiflora* and carvacrol in an experimental model of pulmonary inflammation.

Materials and Methods

Chemicals

OTA, glutathione (GSH), MDA, SOD, catalase (CAT), Thiol, NO and other materials were purchased from Sigma-Aldrich, (St. Louis, MO, USA). Anti-serum against β -actin, BAX, Bcl2, Caspase 3-9-cleaved (Respectively #4970, #2772, #2870, #9664, #7237). In addition, anti-mouse and rabbit horse radish peroxidase labeled IgG (#7076 and #7074, Cell Signaling, USA) were used for check protein levels by western blot method. For Nrf2 pathway analysis (1: 1000) diluted of anti-Keap1, anti-Nrf2, anti-MnSOD, anti-HO-1, anti-toll-like receptor 4 (TLR4), anti-tumor necrosis factor (TNF)- α , interleukine (IL)-1 β , IL-6 and IL-10 were used (Sigma- Aldrich, St. Louis, MO, USA).

Animals and the experimental protocol

In this research 30 male albino mice (weighing 20 to 30 g) were purchased and randomly divided into 5 groups (n=6). Animals were housed in cages at 22 ± 1 °C with temperature conditions of 26-19 26 °C, relative humidity of 70-75%, light intensity of 300 lux, exposure of

12: 12 (light / dark), in additions food, water, and everything else they needed was available in sterile conditions.

Adult mice received a dose of OTA (0.5 μ g/kg) by gavages in 14 day. Group 1: control group normal saline (0.5 μ g/kg); Group 2: OTA (0.5 μ g/kg) negative control group; Group 3: OTA (0.5 μ g/kg) + *Zataria multiflora* (ZM) (25 mg/kg); Group 4: OTA (0.5 μ g/kg) + ZM (50 mg/kg); Group 5: Vitamin E 20 mg/kg + OTA (2 μ g/rat) were treated group.

Biochemical assessment

After treatment, the animals were sacrificed, and their lung tissues were dissected and kept at -80 °C for biochemical analyses. The samples were homogenized in ice-cold normal saline at a volumes of 9 g/L (1: 9 w/v). After centrifuging the homogenates at 4000 rpm/min for 10 min at 40 °C, the supernatant was collected for testing [27]. A spectrophotometer (Jenway 6105 UV/Vis, UK) was used according to the procedures outlined in the assay kits. The broncho alveolar fluid (BALF) supernatants were used for the evaluation of activities of MDA, GSH, SOD, Thiol, NO and CAT [28, 29].

Western blotting

The western blot technique was employed to study molecular damage induced by OTA. Lung tissues from mice were homogenized and the extracted proteins were separated, then transferred from the gel onto a proteins-binding membrane such as nitrocellulose, where their positions are preserved [30, 31]. Antibodies were used as the primary means of detecting specific proteins in the membrane. Protein concentrations were measured in the tissue supernatant with a

protein measurement kit. All separation and detection steps followed kit protocols and referenced other studies [32, 33].

Histopathological analysis

At the end of the treatment period, the animals were sacrificed and their lungs were maintained in 4% paraformaldehyde. Washed lung tissue samples were embedded in paraffin and cut into 5 µm sections for histopathological studies. Hematoxylin and eosin (H&E) as well as periodic acid-Schiff (PAS) staining were used to investigate damage scores [34, 35].

BALF analysis

To examine the BALF, the trachea was first washed using phosphate-buffered saline (PBS), and lavage fluid was recovered. Protein concentration (Beyotime Kit, China) and number of cells (Beckman Coulter) were measured in the collected samples [36].

Wet-to-dry (W/D) ratio assay

To determine the W/D lung weight ratio, lung samples were dried in an oven at 60 °C for 7 days and their weights were recorded [32]. After drying again, the weights were reassessed, and the W/D ratio was calculated [31, 37].

Evaluation of myeloperoxidase (MPO) activity

MPO is a leukocyte-derived enzyme that generates reactive oxidant species, making it useful as an additional indicator of inflammation and tissue damage. MPO activity is quantified based on the reaction between Tetramethyl benzidine (TMB) and H₂O₂, which is read at a wavelength of 650 nm. Neutrophil levels rise substantially during pulmonary inflammation. Pulmonary homogeneous samples are prepared and measured based on the kit protocol [38].

MPO detection followed the manufacturer's guidelines for the mouse MPO Nampox™ MPO activity assay kit.

Statistical analysis: All statistical analyses were performed using one-way ANOVA, followed by Tukey's post hoc test. A P-value lower than 0.05 was considered statistically significant. Partial Eta-squared values were also reported to represent effect sizes.

Results

Extract characterization

Z. multiflora extract, provided by the Pharmacological Research Center of the School of Medicine, was prepared using a Soxhlet extraction system and then analyzed by high-performance liquid chromatography (HPLC) to check the concentration of the active compound, carvacrol. The results of HPLC confirming the presence of carvacrol in the extract are presented in Fig. 1.

Effect of *Z. multiflora* on oxidative stress

As shown in Fig. 2A and F, exposure to OTA (0.5 µg/kg) significantly increased level of MDA and NO as compared to control respectively (P < 0.001). Treatment with *Z. multiflora* (25 and 50 mg/kg) and Vitamin E (20 mg/kg) reduced MDA content in comparison with OTA (0.5 µg/kg) (p<0.01, p < 0.001). In the OTA (0.5 µg/kg) treated group, GSH content was decreased in lungs (Fig. 2B, p<0.01, p<0.001).

Treatment with *Z. multiflora* (25 and 50 mg/kg) and Vitamin E (50 mg/kg) significantly increased GSH content compared to OTA treated rats (p<0.001). Similar results have been obtained from CAT (Fig. 2C, p<0.01-p < 0.001) and SOD (Fig. 2D, p < 0.001 for all cases).

According to results Thiol level as well as CAT and SOD activities were decreased (Fig. 2E, $p < 0.01$ - $p < 0.001$).

Effect of *Z. multiflora* on apoptotic factors (Bax/Bcl-2, Caspase 3 and Caspase 9)

As indicated in Fig. 3A protein expression of Bax/Bcl2 was increased in OTA group ($P < 0.001$). In addition, the protein levels of cleaved caspases 3 and 9 were up-regulated by administration of OTA (0.5 $\mu\text{g}/\text{kg}$). Co-treatment of OTA (0.5 $\mu\text{g}/\text{kg}$) with *Z. multiflora* (25 and 50 mg/kg) or vitamin E significantly decreased the ratio of Bax/Bcl2 in comparison with OTA group ($p < 0.001$ and $P < 0.001$, respectively). Furthermore, administration of *Z. multiflora* (25

and 50 mg/kg) or vitamin E (50 mg/kg) plus OTA inhibited the activation of caspases 3 and 9 (Fig. 3B and C).

Effect of *Z. multiflora* on the structural damage in mice lungs

Structural alternations associated with lung inflammation are shown in Fig. 4. H&E (A) and PAS staining (B) were performed to observe the lung injury score (C), W/D ratio (D), evaluation of total protein content, and (E) lung inflammatory cell counts in BALF from different groups. (F) Lung MPO activity was also determined among the groups (G) ($p < 0.01$ - $p < 0.05$ - $p < 0.001$).




Fig. 1: The results of high-performance liquid chromatography based on the presence of the active ingredient carvacrol in the *Z. multiflora* (50 mg/kg) extract. (C₁₀H₁₄O, 5/1000)




Fig. 2: Effects of *Zataria multiflora* (ZM) extract on Malondialdehyde (MDA) concentrations (A), Glutathione (GSH) (B), Catalase (CAT) activity (C), SOD activity (D), Thiol (E) and NO₂ (F) in the lung tissues mice's. ###p<0.001 vs. control, ***p<0.001 compared to Ochratoxin A group. *Zataria multiflora* resin (Carvacrol). OTA= Ochratoxin-A; Vit E= Vitamin E




Fig. 3: Effects of *Zataria multiflora* (ZM) (25 and 50 mg/kg) on expressions of apoptotic factors following exposure to Ochratoxin-A (OTA) in mice's. The blots and the bars exhibiting the densitometry analysis of western blots for Bax/Bcl-2 (A) ratio, Caspase 3 (B) and Caspase 9 (C) proteins, consecutively. ###P<0.001 vs. control, **P<0.01 and ***P<0.001 compared to OTA-administered mice's.

Discussion

The results of the present study investigated the antioxidant, and anti-inflammatory properties of *Z. multiflora* and its constituent, carvacrol in an experimental model of lung inflammation induced by OTA in mice. Although various studies have documented OTA toxicity in the brain, heart, and kidneys, this study highlights its considerable ability to trigger inflammation, apoptosis, and destruction in lung tissue [39, 40]. Previous studies have revealed that OTA can cross the

blood-brain barrier and exert neurotoxic effects in laboratory rodents [41, 42]. The experimental examined pulmonary toxicity, apoptosis, and OTA-induced inflammation following 14 days of treatment in rats. After the administration period, the animals were examined for weight changes, and the results showed an increase in weight in the samples. The study again elevated pulmonary toxicity, apoptosis, and OTA-induced inflammation after 14 days of gavage administration in

treated rats. Post-treatment examination also revealed weight gain in the samples.




Fig. 4. Effects of *Zataria multiflora* (ZM) in structural damage of lung inflammation. H&E (A) and PAS staining (B) were performed to observe the lung injury score (C), wet-to-dry ratio (D), evaluation of the

Broncho alveolar lavage fluid (E) lung inflammatory cell counts in broncho alveolar fluid from different groups (F) and evaluation of myeloperoxidase (G). OTA= Ochratoxin-A; Vit E= Vitamin E

Pathobiological studies indicate inflammation and tissue destruction in groups that received OTA. The results show the antioxidant and anti-inflammatory effects of *Z. multiflora* receptor groups. Our results indicated that lung levels of MDA and NO metabolites were significantly increased, whereas GSH, CAT and other oxidative-stress-related indicators such as thiol and SOD displayed a significant reduction in the OTA-exposed groups. The results of antioxidant studies align with other research demonstrating that OTA induces oxidative stress in treated specimens [43-45]. According to the present results, apoptosis was particularly observed in toxin-exposed groups. The proteins in this family, especially Bcl-2 and Bax, regulate the apoptotic pathway. By determining the ratio of these two proteins, the rate of cell death can be determined. Activation of these pathways at the cellular level can lead to the formation of apoptosis and eventually the activation of caspases (especially caspase-3, and caspase-9) [46].

Given that Nrf2 is a key pathway in examining the rate of inflammation in various organs and tissues, it was examined the most important factors involved in this pathway. Various studies have shown the effective role of Nrf2 in many diseases caused by oxidative stress [47]. Consequently, it was examined proteins involved in this pathway. Antioxidant enzymes in this pathway such as HO-1, and MnSOD, which are regulated by Nrf2, contribute to

pathogenesis when altered [48]. According to the results of other researchers on the anti-inflammatory effects of *Z. multiflora* [39], the results of this study indicate a decrease in inflammatory factors in the groups receiving *Z. multiflora*. So far, the analgesic and anti-inflammatory effects of the total extract and the essential oil and flavonoid fractions of this plant have been studied. Key inflammatory mediators include TNF- α , IL-1 β , and IL-6. Although the results of this study are similar to those of other studies in terms of the anticancer effects of *Z. multiflora* [49] and the toxic effects of OTA, due to the increase in the level of inflammatory factors as well as OTA appears to cause a specific toxicity in the lung tissue.

According to the results of this study and the toxic effects of OTA and the activation of apoptosis signaling pathways, the protective effects of *Z. multiflora* and its combination with carvacrol has a special ability in an experimental model of lung inflammation.

Conclusions

Our study demonstrated that apoptosis signaling pathways, oxidative stress and inflammation were involved in OTA-induced experimental model of lung inflammation. The present study showed a significant reduction in oxidative stress, inflammation, and apoptosis in the samples receiving *Z. multiflora* through the apoptosis signaling pathway.

Ethical Considerations

The study was approved by the research ethics of Islamic Azad University of Shahrekord, NUM:1402.049.

Funding Statement

The responsible author covered the experiment and other associated costs.

Conflict of Interest

The authors declared no conflict of interest.

Acknowledgment

This manuscript benefited from AI-assisted refinement and formatting support, including

language polishing and abbreviation standardization.

Data Availability Statement

The data presented in this study are available on request from the corresponding author.

Authors' Contributions

F.N. and H.C.M. were responsible for designing the review protocol, conducting the literature review, providing feedback on the manuscripts, writing the manuscript and improving the interpretation of the results. M.S. and E.T. were responsible for writing the manuscript, assembling data, analyzing data, and interpreting analyses.

References

- [1]. Speijers GJA, Speijers MHM. Combined toxic effects of mycotoxins. *Toxicology letters*. 2004; 153(1): 91-8.
- [2]. Kuiper-Goodman T, Scott P. Risk assessment of the mycotoxin ochratoxin A. *Biomedical and environmental sciences* 1989; 2(3): 179-248.
- [3]. Pitt JI. *Penicillium viridicatum*, *Penicillium verrucosum*, and production of ochratoxin A. *Appl Environ Microbiol*. 1987; 53(2): 266-9.
- [4]. Sage L, Krivobok S, Delbos É, Seigle-Murandi F, Creppy EE. Fungal flora and ochratoxin A production in grapes and musts from France. *Journal of Agricultural and Food Chemistry*. 2002; 50(5): 1306-11.
- [5]. Visconti A, De Girolamo A. Fitness for purpose—Ochratoxin A analytical developments. *Food Additives and Contaminants* 2005; 22(s1): 37-44.
- [6]. Varga J, Kozakiewicz Z. Ochratoxin A in grapes and grape-derived products. *Trends in Food Science & Technology* 2006; 17(2): 72-81.
- [7]. Lee TP, Saad B, Khayoon WS, Salleh B. Molecularly imprinted polymer as sorbent in micro-solid phase extraction of ochratoxin A in coffee, grape juice and urine. *Talanta* 2012; 88: 129-35.
- [8]. Abrunhosa L, Paterson RR, Venâncio A. Biodegradation of ochratoxin A for food and feed decontamination. *Toxins* 2010; 2(5): 1078-1099.
- [9]. Duarte SC, Pena A, Lino CM. Human ochratoxin A biomarkers—from exposure to effect. *Critical Reviews In Toxicology* 2011; 41(3): 187-212.
- [10]. Malir F, Ostry V, Novotna E. Toxicity of the mycotoxin ochratoxin A in the light of recent data. *Toxin Reviews* 2013; 32(2): 19-33.
- [11]. Pfohl-Leszkowicz A, Manderville RA. An update on direct genotoxicity as a molecular mechanism of ochratoxin a carcinogenicity. *Chemical Research in Toxicology* 2012; 25(2): 252-62.
- [12]. Malir F, Ostry V, Pfohl-Leszkowicz A, Novotna E. Ochratoxin A: Developmental and reproductive toxicity—An overview. *Birth Defects Research Part B: Developmental and Reproductive Toxicology* 2013; 98(6): 493-502.
- [13]. Minervini F, Giannoccaro A, Nicassio M, Panzarini G, Lacalandra GM. First evidence of placental transfer of ochratoxin A in horses. *Toxins* 2013; 5(1): 84-92.
- [14]. Frizzell C, Verhaegen S, Ropstad E, Elliott C, Connolly L. Endocrine disrupting effects of ochratoxin A at the level of nuclear receptor activation and steroidogenesis. *Toxicology letters* 2013; 217(3): 243-50.
- [15]. Al-Anati L, Petzinger E. Immunotoxic activity of ochratoxin A. *Journal of Veterinary Pharmacology and Therapeutics* 2006; 29(2): 79-90.
- [16]. Pinelli E, El Adlouni C, Pipy B, Quartulli F, Pfohl-Leszkowicz A. Roles of cyclooxygenase and lipoxygenases in ochratoxin A genotoxicity in human epithelial lung cells. *Environmental Toxicology and Pharmacology* 1999; 7(2): 95-107.
- [17]. Xu W, Wang M, Cui G, Li L, Jiao D, Yao B, et al. Astaxanthin Protects OTA-Induced Lung Injury in Mice through the Nrf2/NF-κB Pathway. *Toxins (Basel)* 2019; 11(9).
- [18]. Tao Y, Xie S, Xu F, Liu A, Wang Y, Chen D, et al. Ochratoxin A: Toxicity, oxidative stress and metabolism. *Food and Chemical Toxicology: An International Journal Published for the British Industrial Biological Research Association* 2018; 112: 320-31.
- [19]. Okutan H, Aydin G, Ozcelik N. Protective role of melatonin in ochratoxin a toxicity in rat

- heart and lung. *Journal of Applied Toxicology: An International Journal* 2004; 24(6): 505-12.
- [20]. Han Z, Zhao Z, Shi J, Liao Y, Zhao Z, Zhang D, et al. Combinatorial approach of LC-MS/MS and LC-TOF-MS for uncovering in vivo kinetics and biotransformation of ochratoxin A in rat. *Journal of Chromatography B* 2013; 925: 46-53.
- [21]. Mahboubi M. Management of acute cough by *Z. multiflora* Boiss as an alternative treatment. *Journal of Integrative Medicine* 2018; 16(1): 20-5.
- [22]. Shomali T, Mosleh NZ. *Multiflora*, broiler health and performance: a review. *Iranian journal of veterinary research* 2019; 20(2): 81-8.
- [23]. Khajavi Rad A, Mohebbati R. *Z. multiflora* extract and carvacrol affect cardiotoxicity induced by Adriamycin in rat. *Journal of Basic and Clinical Physiology and Pharmacology* 2018; 30(1): 73-9.
- [24]. Alavinezhad A, Hedayati M, Boskabady MH. The effect of *Z. multiflora* and carvacrol on wheezing, FEV1 and plasma levels of nitrite in asthmatic patients. *Avicenna Journal of Phytomedicine* 2017; 7(6): 531-41.
- [25]. Boskabady M, Jafari Z, Pouraboli I. The effect of carvacrol on muscarinic receptors of guinea-pig tracheal chains. *Phytotherapy Research* 2011; 25(4): 530-35.
- [26]. Richard C, Shabbir W, Ferraro P, Massé C, Berthiaume Y. Alveolar liquid clearance in lung injury: Evaluation of the impairment of the β_2 -adrenergic agonist response in an ischemia-reperfusion lung injury model. *Respiratory Physiology & Neurobiology* 2019; 259: 104-10.
- [27]. Avezov K, Reznick AZ, Aizenbud D. Oxidative stress in the oral cavity: Sources and pathological outcomes. *Respiratory Physiology & Neurobiology* 2015; 209: 91-4.
- [28]. Jammes Y, Steinberg JG, Brégeon F, Delliaux S. The oxidative stress in response to routine incremental cycling exercise in healthy sedentary subjects. *Respiratory Physiology & Neurobiology* 2004; 144(1): 81-90.
- [29]. Wang K, Zu C, Zhang Y, Wang X, Huan X, Wang L. Blocking TG2 attenuates bleomycin-induced pulmonary fibrosis in mice through inhibiting EMT. *Respiratory Physiology & Neurobiology* 2020; 276: 103402.
- [30]. Ji L, Liu R, Zhang XD, Chen HL, Bai H, Wang X, et al. N-acetylcysteine attenuates phosgene-induced acute lung injury via up-regulation of Nrf2 expression. *Inhalation Toxicology* 2010; 22(7): 535-42.
- [31]. Sui H, Luo M, Miao Y, Cheng W, Wen S, Zhao B, et al. Cystic fibrosis transmembrane conductance regulator ameliorates lipopolysaccharide-induced acute lung injury by inhibiting autophagy through PI3K/AKT/mTOR pathway in mice. *Respiratory Physiology & Neurobiology* 2020; 273: 103338.
- [32]. Yu Z, Jiang Y, Sun C. Glucocorticoids inhibits the repair of airway epithelial cells via the activation of wnt pathway. *Respiratory physiology & neurobiology* 2020; 271: 103283.
- [33]. Liu J, Jiang X, Li L, Liu H, Zhang X, Liu K, et al. Iloprost inhibits acute allergic nasal inflammation by GATA3-ILC2 pathway in mice. *Respiratory Physiology & Neurobiology* 2020; 276: 103364.
- [34]. Turner SM, Hoyt AK, ElMallah MK, Falk DJ, Byrne BJ, Fuller DD. Neuropathology in respiratory-related motoneurons in young Pompe (Gaa^{-/-}) mice. *Respiratory Physiology & Neurobiology* 2016; 227: 48-55.
- [35]. Xiang GA, Zhang YD, Su CC, Ma YQ, Li YM, Zhou X, et al. Dynamic changes of mononuclear phagocytes in circulating, pulmonary alveolar and interstitial compartments in a mouse model of experimental silicosis. *Inhalation Toxicology* 2016; 28(9): 393-402.
- [36]. Chen J, Shao Y, Xu G, Lim C, Li J, Xu D, et al. Bone marrow-derived mesenchymal stem cells attenuate phosgene-induced acute lung injury in rats. *Inhalation Toxicology* 2015; 27(5): 254-61.
- [37]. Stenfors N, Bosson J, Helleday R, Behndig AF, Pourazar J, Törnqvist H, et al. Ozone exposure enhances mast-cell inflammation in asthmatic airways despite inhaled corticosteroid therapy. *Inhalation Toxicology* 2010; 22(2): 133-39.
- [38]. Hosseinzadeh H, Ramezani M, Salmani G-a. Antinociceptive, anti-inflammatory and acute toxicity effects of *Z. multiflora* Boiss extracts in mice and rats. *Journal of Ethnopharmacology* 2000; 73(3): 379-85.
- [39]. Limonciel A, Jennings P. A review of the evidence that ochratoxin A is an Nrf2 inhibitor: implications for nephrotoxicity and renal carcinogenicity. *Toxins* 2014; 6(1): 371-9.
- [40]. Ose A, Kusuhara H, Endo C, Tohyama K, Miyajima M, Kitamura S, et al. Functional characterization of mouse organic anion transporting peptide 1a4 in the uptake and efflux of drugs across the blood-brain barrier. *Drug Metabolism and Disposition* 2010; 38(1): 168-76.
- [41]. Kusuhara H, Sugiyama Y. Efflux transport systems for drugs at the blood-brain barrier and blood-cerebrospinal fluid barrier (Part 1). *Drug Discovery Today* 2001; 6(3): 150-6.
- [42]. Petrik J, Žanić-Grubišić T, Barišić K, Pepeljnjak S, Radić B, Ferencić Ž, et al. Apoptosis and oxidative stress induced by ochratoxin A in rat kidney. *Archives of Toxicology* 2003; 77(12): 685-93.

- [43]. Abdel-Wahhab MA, Abdel-Azim SH, El-Nekeety AA. Inula crithmoides extract protects against ochratoxin A-induced oxidative stress, clastogenic and mutagenic alterations in male rats. *Toxicon* 2008; 52(4): 566-73.
- [44]. Pozzo L, Cavallarin L, Antoniazzi S, Guerre P, Biasibetti E, Capucchio MT, et al. Feeding a diet contaminated with ochratoxin A for broiler chickens at the maximum level recommended by the EU for poultry feeds (0.1 mg/kg) 2. Effects on meat quality, oxidative stress, residues and histological traits. *Journal of Animal Physiology and Animal Nutrition* 2013; 97: 23-31.
- [45]. Wei Y, Sinha SC, Levine B. Dual role of JNK1-mediated phosphorylation of Bcl-2 in autophagy and apoptosis regulation. *Autophagy* 2008; 4(7): 949-51.
- [46]. Wang X, Ye Xl, Liu R, Chen HL, Bai H, Liang X, et al. Antioxidant activities of oleanolic acid in vitro: possible role of Nrf2 and MAP kinases. *Chemico-biological Interactions* 2010; 184(3): 328-37.
- [47]. Jeong YH, Park JS, Kim DH, Kim HS. Lonchocarpine increases Nrf2/ARE-mediated antioxidant enzyme expression by modulating AMPK and MAPK signaling in brain astrocytes. *Biomolecules & Therapeutics* 2016; 24(6): 581.
- [48]. Nemati F, Janitermi M, Mohammadpour G. Cytotoxic effect of *Z. multiflora* on gastric cancer cell line (AGS) and normal fibroblast cells. *Cumhuriyet Üniversitesi Fen-Edebiyat Fakültesi Fen Bilimleri Dergisi* 2015; 36(3): 1876-884.