

Letter to Editor

Burn Patients and Augmented Renal Clearance: An Important but Neglected Issue

Farzan Madadizadeh Ph.D., Golnaz Afzal * Ph.D.

Clinical Pharmacy Department, School of Pharmacy, Shahid Sadoughi University of Medical Science, Yazd, Iran

Article history

Received: 2 Jan 2025 Accepted: 1 Jun 2025 Available online: 17 Jun 2025

Keywords

Augmented renal clearance Burn Kidney impairment

Dear Editor,

Patients suffering from major burn injuries represent a unique population of critically ill individuals. An immune and inflammatory response, metabolic changes, and subsequent hypermetabolism accompany them [1]. These conditions can be challenging to manage and may lead to multiple organ failure, affecting various organs such as the heart, liver, and kidneys [2]. The renal failure that occurs in extensively burned patients is typically linked to the failure or dysfunction of other organs, manifesting as multiple organ dysfunction syndrome, which negatively impacts the prognosis [3]. On the other hand, physiological and metabolic changes resulting from burn injuries can impact pharmacokinetic parameters [4]. It can lead to larger volumes of distribution, faster hepatic metabolism, and increased renal clearance [5]. The idea of enhanced renal function and accelerated medication clearance after a major burn injury has been documented as far back as the 1970s [6]. Over time, patients with severe burn injuries have been found to experience an increase in renal clearance, known as augmented renal clearance (ARC). This term, relatively new, typically refers to a creatinine clearance (CrCl) exceeding 130 mL/min/1.73 m² in men or 120 mL/min/1.73 m² in women, as determined through timed urine collection spanning 8 to 24 hours [7].



CC BY-NC 4.0

The phenomenon of accelerated renal clearance is characterized by the rapid elimination of circulating solutes and medications [8]. This can potentially lead to renally cleared antibiotics being at risk below therapeutic concentrations, ultimately putting patients at risk of invasive infections, mortality, treatment failure, and worse clinical outcomes [9]. The prevalence of ARC is estimated to be as high as 70% in critically ill patients, particularly in conditions such as trauma and burns associated with various stress factors [10].

Therefore, understanding and monitoring ARC in burn patients is crucial for optimizing medication dosing and preventing potential complications, ensuring effective and safe pharmacotherapy. Furthermore, the importance of special attention to ARC in burn patients is outlined as follows.

Patients with a larger total body surface area (TBSA), younger age (typically under 50 years), male sex, apparently normal renal function, and lower incidence of organ failure have a well-documented risk of ARC [11, 12].

Currently, there is no ideal substitute indicator or assessment tool available for assessing ARC in burn patients, which poses a significant clinical challenge in detecting and managing this condition [11]. Although measuring urine CrCl is a more accurate method of determination, the measurements are typically limited to the first few days following the burn occurrence [13].

Normal serum creatinine levels may indicate under-recognized ARC, while the use of standard antibiotic dosage regimens is likely to lead to suboptimal concentrations, inferior clinical outcomes, the emergence of multidrug-resistant bacteria, and increased costs for patients and the community [11].

Acute kidney injury occurred in 38% of burn patients, which is more likely to happen in the elderly [14]. The prevalence of ARC was 70.6%, with an incidence of 66.3%, which is more common in younger age groups [15]. Here are two main outcome summaries, as demonstrated in Table 1.

Table 1. Renal dysfunction in burn patients

Type of renal dysfunction	Risk factor	The overall prevalence in the world
Acute kidney injury	Elderly, previous chronic disease [14]	38% [14]
Augmented renal clearance	Young age, apparently normal renal function [11], lower organ failure [12]	70.6% [15]

In essence, while renal injury in burn patients is well-studied, ARC is another crucial aspect that requires attention. Limited data exists on ARC in burn patients; however, it is particularly prevalent in younger individuals, especially males with significant total TBSA involvement. Given

this risk factor, clinicians should be vigilant and consider adjusting dosages upward to improve treatment outcomes. Additionally, healthcare providers must be cautious about the wide spectrum of renal function changes that may occur when caring for burn patients.

References

- [1]. Jeschke MG, van Baar ME, Choudhry MA, Chung KK, Gibran NS, Logsetty S. Burn injury. Nat Rev Dis Primers. 2020; 6(1): 11.
- [2]. Tasleem S, Siddiqui AI, Zuberi MAW, Tariq H, Abdullah M, Hameed A, et al. Mortality patterns and risk factors in burn patients: A cross-sectional study from Pakistan. Burns Open 2024; 8(1): 13-8.
- [3]. Emara SS, Alzaylai AA. Renal failure in burn patients: a review. Ann Burns Fire Disasters. 2013; 26(1): 12-5.
- [4]. Udy AA, Roberts JA, Lipman J, Blot S. The effects of major burn related pathophysiological changes on the pharmacokinetics and pharmacodynamics of drug use: An appraisal utilizing antibiotics. Adv. Drug Deliv Rev. 2018; 123: 65-74.
- [5]. Pruskowski KA. Pharmacokinetics and pharmacodynamics of antimicrobial agents in burn patients. Surg Infect (Larchmt) 2021; 22(1): 77-82.
- [6]. Loirat P, Rohan J, Baillet A, Beaufils F, David R, Chapman A. Increased glomerular filtration rate in patients with major burns and its effect on the pharmacokinetics of tobramycin. N Engl J Med. 1978; 299(17): 915-19.
- [7]. Cook AM, Hatton-Kolpek J. Augmented renal clearance. Pharmacotherapy 2019;39(3):346-54.
- [8]. Shi AX, Qu Q, Zhuang HH, Teng XQ, Xu WX, Liu YP, et al. Individualized antibiotic dosage regimens for patients with augmented renal clearance. Front. Pharmacol. 2023; 14.
- [9]. He J, Yang ZT, Qian X, Zhao B, Mao EQ, Chen EZ, et al. A higher dose of vancomycin is needed in critically ill patients with augmented

- renal clearance. Transl Androl Urol. 2020; 9(5): 2166-171.
- [10]. Xiao Q, Zhang H, Wu X, Qu J, Qin L, Wang C. Augmented renal clearance in severe infections-an important consideration in vancomycin dosing: A narrative review. Front Pharmacol. 2022; 13: 835557.
- [11]. Smeets TJL, Boly CA, Papadopoulos J, Endeman H, Hunfeld NGM. What every intensivist should know about augmented renal clearance (ARC). J Crit Care. 2024; 84: 154541.
- [12]. Torian SC, Wiktor AJ, Roper SE, Laramie KE, Miller MA, Mueller SW. burn injury and augmented renal clearance: A case for optimized piperacillin-tazobactam dosing. J Burn Care Res. 2022; 44(1): 203-206.
- [13]. Mueller SW, Blass B, Davis JN, Deeter LA, Gibson C, Kohler AD, et al. 292 augmented renal clearance in burn patients: Incidence and discordance with standard creatinine clearance estimation. J Burn Care Res. 2023; 44(S2): 187-88.
- [14]. Folkestad T, Brurberg KG, Nordhuus KM, Tveiten CK, Guttormsen AB, Os I, et al. Acute kidney injury in burn patients admitted to the intensive care unit: A systematic review and meta-analysis. Crit Care. 2020; 24(1): 2.
- [15]. Mueller SW, Blass B, Molina KC, Gibson C, Krsak M, Kohler AD, et al. Augmented renal function in burn patients: Occurrence and discordance with commonly used methods to assess renal function. J Burn Care Res. 2023; 44(6): 1298-303.